

trifoliate, petioled cauline leaves, has been generally taken in Canada to be typical *S. Canadensis*, L.

*OSMORRHIZA BREVISTYLIS*, DC.

From Prince Edward Island west to Lake Winnipeg.

*OSMORRHIZA LONGISTYLIS*, DC.

From Nova Scotia west to the Saskatchewan.

*OSMORRHIZA NUDA*, Torr.

From the Eastern slope of the Rocky Mountains west to Vancouver Island.

*OSMORRHIZA OCCIDENTALIS*, Torr.

Mountain woods at Ainsworth, Kootanie Lake, B.C., alt. 5,000 ft., 1890. (*John Macoun*.) A new station for this plant.

*CICUTA CALIFORNICA*, Gray.

New Westminster, B.C.; Ainsworth, Kootanie Lake B.C. (*John Macoun*.) Not before recorded from British Columbia mainland.

*CARUM CARVI*, L.

Waste places near the brick-yard at Banff, Rocky Mountains. (*John Macoun*.)

*LIGUSTICUM GRAYI*, C. & R.: *Macoun. Cat. Can. Plants*, Vol. II., p. 327.

Woods on the mountains at Ainsworth, Kootanie Lake, B.C., alt. 5,000 feet. (*John Macoun*.)

*LIGUSTICUM SCOPULORUM*, Gray.

Specimens collected by Prof. John Macoun, at Roger's Pass, Selkirk Mountains, B.C., in 1890, have been doubt-