Diarrhœa, 27, 70, 71, 199, 200 fruit, to avoid, in, 51 flour ball, recommended, 43 Diet, after two years old, 52-60 and hair, 176 errors in babe's, 17 for school children, 61-67 in colds, 75, 76 in constipation, 68, 69 in diarrhœa, 70 in eczema, 76 in fever, 77, 78 in illness, 68-80 in indigestion, 71-73 in rickets, 73—75 in rheumatism, 279 in tuberculosis, 78-80 Digestion and rest, 75 and skin diseases, 76 barley food, to help, 30 eggs and, 46, 47 food and feeding, 28 infant's, 19, 20, 23, 39 starchy food and, 41 Dinner, school children, 65, 66 Diphtheria, 208, 262-265 Discharges, the, 198 from ears, 170 from eyes, 152, 158 from nose, 75, 258 involuntary, 251, 252 in ailing child, 245 in diarrhœa, 199 in typhoid fever, 272 Diseases, children's, 244—286 Disinfectants, 253, 254 Disinfection, when necessary, 153, 257, 259, 268, 272, 273, 284 Disorders in young children, 17,23 incident to teething, 146-148

Distended stomach, infan, a 44 Drainage, when bad, 280 Drawers, 104 Drowning, 304, 305 Drum, ear, 164 EARACHE, 167 Ears, 164-170 Eating too fast, 72 Eczema, 76, 211 Eggs, 46, 47, 53 at meals, 64-67 fried, indigestible, 47 in constipation, to avoid, 69 Electrolysis and hair, 179 Emergencies, 287-306 Enemata, nutritive, 247-249 Enlarged glands, 238, 239 tonsils, 239, 240 Epilepsy, 299 Erythema, 76, 210 Eustachian tube, 164 Evacuations, 198-200 See under AILMENTS. Excitable children, 217 Exercise, 277, 285 in menstruation, 319 Expectoration in consumption, 283 External piles, 237, 238 Extra fingers, 233 Extraction of teeth, 151 Eyes, 152-163 FAINTING, 299 Farina, 42 Farinose, 45 Fat, 28, 281 condensed milk makes, 35 Fats, and heat and energy, 6; in food, 62, 79