

A single specimen received from Mrs. Annie Trumbull Slosson. Readily separated from the described American species by the colour of the pleuræ.

*Tenthredo bilineatus*, n. sp. ♀.—Black, with the following parts yellow: the clypeus, the labrum, the mandibles except at apex, a spot on the front beneath the antennæ, an ovate spot on the antennal ridges above the base of each antenna, the lower half of the cheeks, an elongate mark on the inner margin of the eye (this mark is half as long as the inner margin of the eye, reaching the meso-caudal angles of the eye, obliquely truncated in front, roundly emarginate on its mesal side, extending slightly beyond the caudal margin of the eye, a fine spur extending from the middle of its caudal margin to an elongate, quadrangular spot along the latero-caudal margin of the head, sub-interrupted from the mark on the cheeks, not extending mesad beyond the mesal margin of the spot or the inner margin of the eye), a small spot on the vertex, in a line with the spots on the latero-caudal margin of the head and caudad of the ocelli, the tegulæ, the collar, two lines on the mesonotum, converging behind, a broad line from the scutellum to the base of the anterior wings, the cenchri, a small triangular spot on the cephalo-dorsal corner of the mesopleuræ, a spot above the posterior coxæ, the basal membrane, two large spots on the sides of the basal plates, the trochanters, the bases of the femora, the front tibiæ and tarsi, and the middle tibiæ slightly beneath; the following parts rufous: the first segments of the antennæ entirely and the second and third on the inner side, the front and middle femora, the tibiæ above, the posterior femora beneath at apex, the posterior tibiæ, the middle and posterior tarsi, the venter, a narrow margin to the tergal segments one to three, and the tergal segments beyond the third; the third segments of the antennæ one-third longer than the fourth; the clypeus emarginate; the wings hyaline, slightly yellowish; the veins black; the costa and the base of the stigma luteous. Length. 6 mm.

Habitat—Ithaca, New York.

This species will be readily recognized by the markings on the head and mesonotum and the colour of the basal segments of the antennæ and the apex of the abdomen.

*Tenthredo pallipunctus*, n. sp. ♀.—Black, with the following parts yellow: the clypeus, the labrum, the mandibles except at apex, the lower part of the cheeks, a line on the collar, a spot above the posterior