

then directed the treatment to the ulceration of the stomach, and gave iodine 3d, at night, conium in the morning, and arsenicum two or three times through the day, according to the amount of vomiting. This was continued until the vomiting of the dark substance wholly ceased, two months after the first change, and a transparent mucus was discharged. I then gave merc. sol. 3d, at night, continued the conium in the morning, and Lycopo. in the afternoon, for two or three weeks, and until the discharge of this character of mucus ceased. I then gave Cal. Carb. and Coni. alternately, at intervals of 24 and 48 hours, until there was entire reaction of the organism and the enlargement of the thyroid gland disappeared. The patient has remained well since, seven years.

Case 2nd. Mrs. E., aged 22, sanguine, lymphatic temp., full habit — was attacked with convulsions, which continued for thirty-six hours—the patient wholly unconscious. Gave Nux. Hyos. and camphor, alternately, until the convulsions ceased—upon which, a fever, with severe pain extending from the stomach throughout the œsophagus, with tumors, occurred, apparently from the centre of the sternum through the cardiac orifice; restlessness, with constant nausea accompanied with vomiting of white slimy mucus; pulse 120 to 130, tense and twitching. Gave Bell. and Merc. viv. alternately, every three hours, until the fever, restlessness, and vomiting abated, and deglutition became more tolerable. She then commenced ejecting a substance like dark coffee grounds, mixed with mucus, from the stomach, sometimes quite large quantities. I immediately gave Iodine 3d, and alternated with it Arsenicum and Conium, every four hours, until in the course of ten days the ulcers broke and discharged a large quantity of bloody pus, which continued for several days, gradually lessening as the ulcerated cavities healed; extending the time between the doses, as the case continued to grow better. This case was well in two months' time, and she has had no return of the disease since. three years and a half.

Case 3d. Mrs. W., aged 22. Called 8 P. M., to see her; had been confined three days before; first child; child large and healthy. Patient rather large, full habit and sanguine temp.; had severe pains through the whole abdomen, which were very tense, extremely tender to the touch, and very hot; was attacked with chills about 4 P. M.; pulse full, tense, and frequent. Gave Acon. in solution 3d, and Bell & Puls (as the Lochia was entirely suppressed,) in alternate doses every three hours. In forty-eight hours patient well and remained so.

Case 4th. Mrs. M., age 25, was confined with second child; doing well until the fourth day, in the afternoon, was taken with severe chills.