

- BLACK AND WHITE WARBLER, *Mniotilta varia*—Probably the most numerous warbler.
- NASHVILLE WARBLER, *Vermivora rubricapilla*—A few seen each year.
- TENNESSEE WARBLER, *Vermivora peregrina*—August 21st, 1914, one seen.
- PARULA WARBLER, *Compsothlypis americana*—A number seen each year.
- CAPE MAY WARBLER, *Dendroica tigrina*—1914 the first year this bird was recorded. Twelve seen in one day.
- YELLOW WARBLER, *Dendroica aestiva*—A few seem to breed in this locality, but not at all common.
- BLACK-THROATED BLUE WARBLER, *Dendroica caerulescens*—Also breeds in small numbers.
- MYRTLE WARBLER, *Dendroica coronata*—Common.
- MAGNOLIA WARBLER, *Dendroica magnolia*—Quite a common migrant.
- CHESTNUT-SIDED WARBLER, *Dendroica pensylvanica*—Common.
- BAY-BREASTED WARBLER, *Dendroica castanea*—Common during migrations.
- BLACKPOOL WARBLER, *Dendroica striata*—August 21st, 1914, one seen.
- BLACKBURNIAN WARBLER, *Dendroica fusca*—Very common during migrations.
- BLACK-THROATED GREEN WARBLER, *Dendroica virens*—Very common.
- OVEN-BIRD—*Seiurus aurocapillus*—Common.
- WATER-THRUSH, *Seiurus noveboracensis*—Numbers breed in the swampy places. In August they may be seen feeding in the open along the shore among the drift wood and other debris.
- MOURNING WARBLER, *Oporornis philadelphia*—Keeps to the swampy thickets and rarely seen in August.
- MARYLAND YELLOWTHROAT, *Geothlypis trichas*—Quite a common bird.
- WILSON WARBLER, *Wilsonia pusilla*—August 19th, 1914, adult male seen. Only record.
- CANADIAN WARBLER, *Wilsonia canadensis*—Common, especially during migrations.
- AMERICAN REDSTART, *Setophaga ruticilla*—Very common.
- CATBIRD, *Dumetella carolinensis*—Common.
- HOUSE WREN, *Troglodytes aedon*—Fairly common.