The species are :-

Omus dejeanii, Reiche. B. C., Van. Isl. A large back insect, 18 to 20 mm. in length, the elytra marked with rather deep, irregular foveæ.

O. Audouinii, Reiche. B. C., Van. Isl. Resembles the preceding, but the size is less (13 to 18 mm.) and the elytra simply irregularly punctate. The species of Omus do not resemble Cicindela in appearance, but approach rather the Carabidæ, from which they may be easily separated by the antennal characters.

Cicindela longilabris, var. montana, Lec. Resembles longilabris proper, but is black, shining, smoother, the markings absent or only a slender median band present. Has been found in the Northwest Territory. J. B. Tyrrell (teste Harrington).

C. scutellaris, var. unicolor, Dej. Like Lecontei, but green or blue, without marks of any kind. Taken by J. B. Tyrrell in the Northwest Territory.

C. sexguttata, var. patruela, Dej B. C., Rev. Geo. W. Taylor. This is a variety of sexguttata in which the middle band is more or less distinct.

C. purpurea, var. decem-notata, Say. This is a form in which the middle band is more deflexed than in typical purpurea. The elytra are ordinarily ornamented also with four white dots. B. C. (teste Leconte).

C. formosa, Say. Fort McLeod (teste Fletcher), also in N. W. T. This is like *generosa*, but the ground colour is reddish cupreous instead of brown.

C. hyperborea, Lec. Methy Portage, H. B. T. (teste Leconte). I have never seen this species. It is rather small (12 mm.) reddish cupreous with markings somewhat resembling *vulgaris*, but the hinder portion of the median band is straight instead of being curved.

C. vulgaris, var. vibex, Horn. Vanc. Isl., Rev. G. W. Taylor (teste Harrington). A green variety of vulgaris in which the median band does not reach the margin.

*C. oregona*, Lec. A variety of *repanda*, brownish-bronze or olive, middle band not dilated on the margin. White dots large, lunules interrupted. B. C., Van. Isl.

C. pusilla, Say. Hudson's Bay Terr. (teste Schaupp). A small black species, (12 mm.) blue or greenish beneath, with tibiæ pale at base, labrum three-toothed. Elytra with faint basal and apical lunule and recurved median band extended on margin. All but the marginal part of this band is sometimes wanting. Unknown to me.