

flushings of the face from which he had suffered for several weeks. On his way to my house he heard voices apparently saying to him "What is the use of your going to a physician? You are of no use in the world. Go and jump into the river. Jump off the ferryboat; jump, jump, now; at this very instant," and so on. He stated that it was impossible for him to follow his profession, for that the voices interfered to the extent of preventing his clearly distinguishing what was being said in his presence. Even as he was talking to me the hallucinations of hearing were present in full force.

These voices did not actually impose upon his intellect, but he stated that he was conscious of a gradually increasing inability to resist accepting them as realities.

Although there were many of the symptoms of cerebral hyperæmia present, I was induced from the fact that the disorder had come on immediately after bathing in the ocean, during which water had entered the ears, to examine these organs in the very beginning of my interview. Both ears were found full of inspissated cerumen. This was thoroughly softened by the solution of soda in glycerine, and removed by syringing with warm water. On the instant the voices ceased and the patient left, feeling as he said entire relief from his annoying symptoms.

I heard no more of this patient till about two months afterwards, I read in the newspapers of the day that he had been violently abusive in court of the judge on the bench, and had been punished by fine and imprisonment for contempt, and soon afterward his wife called to tell me of the trouble into which her husband had gotten. As she explained it to me he had imagined that the judge was calling him names and cursing him, and had replied in like manner. I had no doubt that there was an accumulation of cerumen, and that the hallucinations of hearing had returned in so aggravated a form as to convince the intellect of their reality. A letter from me to the judge secured his release, and on his visiting me I found my suspicions confirmed. The impacted cerumen was removed, and so far as I know there has been no recurrence of the disorder.

These are only a part of the instances in which impacted cerumen has caused cerebral symptoms that have fallen under my notice, but they are typical, and nothing would be gained by detailing the others.

As regards the cause of noises in the ears I have no information to offer except to state that it is not the mere stoppage of the external meatus by impacted cerumen, for such closure does not give rise to any subjective sensation. It is true that if the canal be stopped by the finger a sound is heard, but this is derived entirely from the body, and is probably from the action of the heart, the circulation of the blood through

the tissues, muscular contraction, etc. A cork or other substance put into the ear so as to close the canal and left there without being held by the hand does not give rise to any sound. If, however, the fingers hold it in place, it transmits the sound from them as would any other solid substance.—*N. Y. Hosp. Gazette.*

LONDON LETTER.

Perhaps the most interesting communication made to any of our societies lately is that of Dr. Matthews Duncan to the Medical Society, on *Antiseptic Midwifery*. So important was it, and listened to with every attention by a distinguished audience, that an abstract of it may be acceptable to your readers. Being a great personal friend of Prof. Lister's, having left the northern metropolis at nearly the exact time Prof. Lister turned his steps southward, it might *a priori* be surmised that Dr. Duncan would be an advocate of the antiseptic plan of treatment. Consequently a large number of practitioners came to hear, and also to learn how antiseptics are applied to every-day midwifery. Dr. Duncan commenced by saying that there is no subject which excites more professional interest or more interest among the general public than that of puerperal deaths. A wife, the mistress of a household, the solace of her husband, the proud mother of a number of happy children, is suddenly snatched away after an auspicious event. There is something so sad about such deaths that all would welcome with heartfelt joy any plan which promises to lessen such disastrous events. Puerperal deaths own various causes, but by far the most frequent and prevalent causes are septicæmia and pyæmia. Both these diseases involve or imply inflammatory processes, and both are essentially septic. It is against them that antiseptic midwifery wages war, and in which, he said, it had already achieved great success. The object of the paper was to spread and diffuse further knowledge on this important matter, and to stimulate further inquiry into it, with a view to the more general adoption of the beneficent antiseptic methods. Already, said Dr. Duncan, more pain is prevented, more life saved by antiseptic methods than by all the recent improvements of modern midwifery combined; and there is no prospect half so bright and encouraging as that held out by the general adoption of the antiseptic treatment of the parturient condition. And, it is certain, all fervently wish that these high hopes may be realized. He would not, he said, proceed to discuss that division of the subject, the treatment of the blood by which the fermentation or sepsis is carried throughout the organism, as by the use of hyposulphites, introduced by Polli, of Milan. He would