

*P. MAXIMUM*, *Jacq.* (*P. jumentorum*, *Pers.*—*P. trichocondylum*, *Steud.*—*P. fasciculatum*, *Pl. Carib.*) “Guinea grass.” Hab. Tropical Africa.

*ZEa MAYS*, *Kth.* “Indian Corn.” Hab. unknown, supposed to be American.

*ZEa JAPONICA.*

*AVENA SATIVA*, *L.* “Oat.” Grows well for a time and then dies off before ripening seed. Oats are generally sown in ground intended for a potato crop, and when about a foot in height, are dug under in order to manure the ground for the coming crop.

*TRITICUM VULGARE*, *L.* “Wheat.” Grows well in some places, and produces fair grain. In former years it was more extensively cultivated, and bread was frequently made in farm houses, but of late years its cultivation has ceased altogether.

*HORDEUM VULGARE*, *L.* “Barley.” Grows well and ripens, but is seldom cultivated as a crop.

*SACCHARUM OFFICINARUM*, *L.* (*S. violaceum*, *Tuss.*) “Sugar Cane.” Grows well, especially in low moist ground.

*SPOROBOLUS INDICUS*, *R. Br.* (*Agrostis*, *L.*—*S. tenuissimus*, *P. B.*—*S. elongatus*, *R. Br.*) Hab. Warmer regions of both hemispheres.

*CHLORIS PETRACA*, *Thunb.* (*Eustachys*, *Desv.*—*C. Swartzii et septentrionalis*, *C. Müll.*)

*CYNODON DACTYLON*, *Pers.* This grass is widely distributed over the tropical and temperate regions of the globe.

*PASPALUM DISTICHUM*, *L.* (*P. litorale*, *R. Br.*—*Digitaria paspaloides*, *Dub.*)

*P. FILIFORME*, *Sw.* (*P. Swartzianum*, *Flugg.*) Hab. West Indies.

*STENOTAPHRUM AMERICANUM*, *Schrk.* (*Rottbællia dimidiata*, *Sw.*—*R. stonolifera*, *Poir.*—*Diastemanthe platystachys*, *Steud.*) Hab. North and South America.

*ANDROPOGON SCHÆNANTHUS.* (*Cymbopogon Schænanthus*, *R.* et *S.*) Hab. East Indies.

*PHALARIS CANARIENSIS*, *Kth.* Hab. Europe.