orth-West

Time

of Seeding.

Aug.-Sept.

July-Sept. July-Aug.

July-Aug.

uly-Sopt. ummer ..

ıly

pt.

ly-Oct...

ly-Sept.

g.-Sept.

ly-Oct.

Territories with their chief characters. - Continued.

Colour, Size,
Arrangement of Flowers
and other
Characters of the Plant. Method of Propagation and Distribution. Place of Growth Methods Eradication. Products Injured. Whole plant dark green, the Seeds, floods..... Grain fields, par numerous very small flowers in a tall wand-like, leafy paniole.

Lilac; ¾ in.; running rootsatocks, atocks, Lilac purple, 2 in.; whole plant grayish.

Grain fields, par ticularly on stubble, hay, market value of land.

Fields, grain, pastures, all crops.

Coasidos, summer fallow.

Bright fall and spring, summer fallow.

Fields, grain, pastures, all crops.

Coasidos, summer fallow.

Summer fallow. Pink 1/4 in., solitary; exuding Seeds, running Grain fields....... Summer fallow, cultimited branched, almost rootstocks. much branched, almost leafless.
Blue; ¾ in., few; loose panicle; glaucous.

Pale yellow; ¼ in.; corymb; leaves heart-shaped at base with many soft spines and two sharp suricles.
Pale yellow; ¼ in. corymb; leaves less divided, more prickly, the auticles at the base rounded.

**Total Corymb; leaves less divided, more prickly, the auticles at the base rounded.

**Total Corymb; leaves less divided, more prickly, the auticles at the base rounded.

**Total Corymb; leaves less divided, more prickly, the auticles at the base rounded.

**Total Corymb; leaves less divided, more prickly, the auticles at the base rounded. Piuk, § in..... Seeds, rootstocks. Mead o wa, wet fields on alkaline lands. Rich soil, all crops Mow whils in bloom, plough, hoed crops. Blue, 16 in., axillary, on leafy Seeds carried by Grain fields, road-racemes. Summer fallow, cul-racemes. Pink or white, 2 in. solitary. Seeds, running Fields........... Cultivate frequently. root stocks. White or lilao, 14 in., umbel-Seeds...... Fields, gardens, Cultivate, hoe. like clusters.