

of ocelli terminating behind in a deep puncture ; clypeus short, scarcely emarginate ; palpi piceous ; antennæ long, stout ; joints 3 to 9 sub-equal, terminal joint blunt at apex.

Thorax polished ; tegulæ, with a large angulated spot before on prothorax, pale rufous ; legs rufous, except base of coxæ and tips of some of the joints of tarsi, which are more or less blackish ; wings hyaline, iridescent ; nervures and stigma black, a spot in centre of second sub-marginal cell, lanceolate cell without crossline, two middle cells in posterior wings.

Abdomen long, flattened, sides parallel ; segments 2 to 5 of tergum with the apical margin narrowly pale rufous.

Described from two specimens captured in May.

6. *TENTHREDO SEMICORNIS*.—♂. Black, abdomen and legs partly rufous ; length 0.40 inch ; expanse of wings 0.85 inch.

Head wider than thorax, excavated in front, with a strong ridge above each antenna ; clypeus, labrum and base of mandibles white ; antennæ moderately stout and about as long as the abdomen ; five basal joints black (the third with a narrow rufous ring at base), four terminal joints white, except the extreme apex of last which is black.

Thorax black, except a white spot on flanks, a larger one above the posterior coxæ, and another on lateral margins of the basal plates ; wings hyaline, stigma and nervures blackish ; tegulæ and base of costa rufous ; legs ferrugineous, anterior pair paler ; coxæ and trochanters partly white with base black ; a line on femora above, a spot at apex of tibiæ within, and posterior tarsi, except last joint and base of first, black.

Abdomen black at base and apex ; segments 3 to 6 rufous, margined with black ; ventre rufous except two apical segments.

Described from a single specimen captured in the city on 9th June, 1886, by Mr. J. A. Guignard. This species has the appearance of *rufopediba* Nort., with the antennæ of *grandis* Nort., by which it may be easily recognized.

7. *TENTHREDOPSIS EVANSII*.—♀. Yellow, with black and green markings ; length 0.45 inch ; breadth of wings 0.90 inch.

Head yellow, except an oval black spot surrounding the ocelli, within this black patch are two short yellow lines, one on each side of lower ocellus ; clypeus squarely emarginate, pubescent, as also labrum which is margined with green ; mandible yellow at base, centre green and tip