

BAYER'S MACEUTICAL PRODUCTS

(Trade Mar.)

MATOSE A tasteless, odourless, nutrient meat it contains an the arcumulation of the meat in an easily soluble for has been extensively employed be of the greatest service in and i. consun tion, diseases of the stomach and intestinal tract, chlorosis and rickets. It is of great value in convalescence from all diseases. SOMATOSE strengthens the muscles and stimulates the appetite in a remarkable manner. Dose for adults: a level teaspoonful three to four times a day with milk, gruel, coffee, etc.

IRON SOMATOSE (Ferro-Somatose). A first-class tonic, containing the albuminous substances of the meat (albumoses) organically combined with iron. Special indications: Chlorosis and Anaemia. Daily dose: 75 to 150 grains.

MILK-SOMATOSE (Lacto-Somatose). A strength giving food containing the albuminous matter (albumoses) of the milk.

TRIONAL A most reliable and quickly-acting hypnotic of the Sulfonal group. Dose: 16 to 20 grains, in a large cup of hot liquid,

IODOTHYRINE The active principle of the thyroid gland. It is most efficacious in Strumous Diseases, Myxoedema, Obesity, Rickets, Psoriasis, Eczema, and Uterine Haemorrhages. Dose : grains two to eight times a day for adults; 5 grains one to three times daily for children.

LYCETOL

RHENACETINE-BAYER PIRERAZINE-BAYER

Solvent. Mas a marked effect on the diuresis. Dose: 16 to 32 grains daily.

An Iodine Cicatrisant which is an excellent odourless substitute for Iodoform and highly recommended for Burns, Wounds, Scrofulous Ulcerations, etc.

EUROPHEN A perfect substitute for Iodoform. Odourless and non-toxic. Has a covering power five times greater than Iodoform. Especially useful in Ulcus molle et durum,

LOSOPHAN cresoltriiodide particularly efficacious in the treatment of all kinds of cutaneous disorders caused by animal parasites.

A new silver pre-paration. Most **PROTARGOL** reliable in cases of Gonorrhoea. Antiseptic wound healer. Excellent results cases of Gonorrhoeal Ophtalmia. Solutions of 1/4 to 2%. Ointments.

TANNIGEN An almost tasteless intestinal astringent. Most efficacious in Chronic, Acute and Summer Diarrhoeas, Adult dose: 8 grains every three hours.

TANNOPINE (Formerly "Tannone "). new intestinal astringent. Special indications: Tuberculous and non-tuberculous Enteritis, Typhus. Dose: 15 grains, three or four times daily.

SALOPHEN Specific for Influenza, Headache, Migraine, Acute Articular Rheumatism, Chorea, Tartrate of Piperazine Sciatica. Dose: 15 grains four to six Anti-Arthritic, Uric times daily. In powders, etc.

SULFONAL BAYER SALOL-BAYER

Samples and literature may be had on application to the

pot in Canada for all "BAYER'S"

WHOLESALE ONLY