

PAPAVERACEÆ.

ARGEMONE MEXICANA, *L.* "Lady Thistle." Very common in STERIP
gardens and waste ground which has been cultivated. From its
flowers is sometimes made a yellow dye, with which the islanders
colour ribbons and other small articles. It flowers about the RESEDA
end of March or beginning of April. The plant appears to have Egy
a very wide geographical range, being found in the northern as
well as southern States of America, West Indies, India, China, ELACOU
and probably all over the tropical as well as temperate regions
of the globe.

PAPAVER — ? Varieties cultivated.

FUMARIACEÆ.

FUMARIA OFFICINALIS, *L.* Hab. Europe.

CRUCIFERÆ.

LEPIDIUM VIRGINICUM, *L.* "Pepper Grass." A common weed
in waste ground.

IBERIS VIOLACEA, *D. C.*

COCHLEARIA OFFICINALIS, *L.* "Scurvy Grass." Common every PORTU
where along the shore, sometimes attaining a large size, almost All
a bush, in sheltered places beneath the cliffs of the south shore gene
at Devonshire Bay. In flower March and April. It is used as and
a cure for diarrhœ, and also for cleansing the blood. boile
are f

C. ARMORACEA. "Horse Radish." Cultivated.

NASTURTIUM — ? Varieties cultivated.

CHEIRANTHUS — ? "Wall Flower." Varieties cultivated.

MATTHIOLA — ? "Stock." Wild among the rocks below SESUVI
Gibb's Hill light house, south shore of Port Royal. Varieties mud
also cultivated in gardens. pera
MESEM
vated

BRASSICA — ? "Cabbage." Varieties cultivated.

BETA — ? "Beet." Varieties cultivated.

SINAPIS — ? "Mustard." A very troublesome weed growing
in cultivated ground. I think there are two species.

RHAPHANUS — ? "Radish." Varieties cultivated.

MALCOMIA MARITIMA, *D. C.* Hab. S. Europe.

MALVA
MODIO
Mal