

CHICKENPOX VACCINE A disease that can be very serious for children and adults who have not had it before. It can cause a fever, a rash, and sometimes complications such as pneumonia or encephalitis.

In December 1988, the original chickenpox vaccine had been replaced by the new varicella-zoster protein. This vaccine is more effective and safer than the old one. It has been recommended for all children and teenagers. It is also safe for pregnant women and for people with weakened immune systems.

The new chickenpox vaccine is given in two doses. The first dose is given at age 12 months. The second dose is given at age 4 years. It is given to children who have not had chickenpox before. It is also given to children who have had chickenpox but did not have a good response to the first dose. It is also given to children who have had chickenpox but did not have a good response to the first dose.

The New Zealand Department of Health's approach to chickenpox vaccination against the virus. The extension of the program to include all children aged 12 months and older is due to the high rates of infection among children and teenagers. The new vaccine is safe and effective. It is also safe for pregnant women and for people with weakened immune systems.

The New Zealand Department of Health's approach to chickenpox vaccination against the virus. The extension of the program to include all children aged 12 months and older is due to the high rates of infection among children and teenagers. The new vaccine is safe and effective. It is also safe for pregnant women and for people with weakened immune systems.

The New Zealand Department of Health's approach to chickenpox vaccination against the virus. The extension of the program to include all children aged 12 months and older is due to the high rates of infection among children and teenagers. The new vaccine is safe and effective. It is also safe for pregnant women and for people with weakened immune systems.

The New Zealand Department of Health's approach to chickenpox vaccination against the virus. The extension of the program to include all children aged 12 months and older is due to the high rates of infection among children and teenagers. The new vaccine is safe and effective. It is also safe for pregnant women and for people with weakened immune systems.

The New Zealand Department of Health's approach to chickenpox vaccination against the virus. The extension of the program to include all children aged 12 months and older is due to the high rates of infection among children and teenagers. The new vaccine is safe and effective. It is also safe for pregnant women and for people with weakened immune systems.

The New Zealand Department of Health's approach to chickenpox vaccination against the virus. The extension of the program to include all children aged 12 months and older is due to the high rates of infection among children and teenagers. The new vaccine is safe and effective. It is also safe for pregnant women and for people with weakened immune systems.

The New Zealand Department of Health's approach to chickenpox vaccination against the virus. The extension of the program to include all children aged 12 months and older is due to the high rates of infection among children and teenagers. The new vaccine is safe and effective. It is also safe for pregnant women and for people with weakened immune systems.