

and a-half lines. The transverse diaphragms are not visible in the specimens examined. The walls of the separate corallites are thick and concentrically wrinkled.

One specimen with corallites two lines in diameter appears to be a variety of this species.

Formation and Locality.—Trenton. Blue Point, Lake St John.

Collector.—J. Richardson.

Genus PALÆOPHYLLUM (Billings).

Generic characters.—Corallum fasciculate or aggregate; corallites surrounded by a thick wall; radiating septa extending the whole length; transverse diaphragms either none or rudimentary; increase by lateral budding.

This genus only differs from *Petraia* or *Streptelasma* by forming long fasciculate or aggregate masses instead of being simple.

PALÆOPHYLLUM RUGOSUM (Billings).

Description.—Corallum in large aggregations of scarcely separate corallites, which where they open out upon the surface of the rock are from one to six lines in diameter, the average adult size being about four lines. Radiating septa reaching the centre; about twenty-two septa in a corallite four lines in diameter, with an equal number in a rudimentary state between.

The great disparity in the size of the tubes in the same mass is owing to the mode of increase and gradual growth of the young corallites. These, of all sizes from one line in diameter and upwards, are uniformly intermingled with the adult individuals.

Formation and Locality.—Trenton. Lake St. John, Little Discharge.

Collector.—J. Richardson.

PETRAIA RUSTICA (Billings).

Description.—Straight or slightly curved, covered with a strong epitheca, which is more or less annulated with broad shallow undulations; radiating septa about one hundred or usually a little more; much confused in the centre, where they form a vesicular mass; every alternate septum much smaller than the others, only half the whole number reaching the centre. Length from two inches and a half to three inches and a half. Diameter of cup one inch to one inch and a half; depth of cup half an inch or somewhat more.