

tion of these phenocrysts of labradorite would produce the so called Huronite. The fine-grained portion of the rock in which these crystals have been developed is a fresh aggregate composed chiefly of plagioclase (labradorite), augite and olivine. The ophitic or diabasic structure is very pronounced. The plagioclase is usually idiomorphic forming an interlacing network of lath-shaped crystals, the interstices of which are filled with augite and olivine. The augite possesses the reddish colour and pleochroism so common in diabase, the larger grains showing frequent distortion and occasional dislocation. Both the feldspar and augite exhibit undulatory extinction as an effect of pressure. The olivine, as usual, occurs in irregular, more or less rounded individuals, only very rarely presenting sharp crystallographic outline. Commonly, it is rather fresh, showing a colourless or light greenish section with characteristic high relief, rough surface and brilliant interference colours. It is rarely so fresh, however, as to be without traversing fissures filled with more or less opaque alteration products. In many instances the original olivine grain is represented by a greenish or yellowish material, probably serpentine. Small scales or grains of opaque iron ore (magnetite) are associated with this serpentine indicating that they were also a result of the decomposition of the olivine. Less frequently, perhaps, the olivine shows a very interesting and rather unusual alteration to talc, but the resulting scales of this mineral were so small that this could not be ascertained beyond dispute. The talc is of a very pale green colour, slightly pleochroic, and exhibits very brilliant interference colours between crossed nicols. It occurs as a matted or felted aggregate of very minute scales filling the original olivine grain. The talc is usually accompanied by more or less opaque iron ore and occasionally some chlorite. (1) A considerable quantity of biotite is present which in some cases has undergone considerable "bleaching" owing to the removal of iron, while in other cases it is altered to chlorite. Apatite is also a tolerably abundant accessory constituent. The magnetite occurs usually in irregular black grains, most of which have resulted from the decomposi-

---

(1) Vol. III. Geol., Wisconsin, p. 235.