

Expanse 18 mil. New York, L. F. Harvey, July 27 ; Texas, Belfrage (without number, among *communis*), Sept. 25.

Botis flavidissimalis, n. s.

Size moderate. Entirely bright deep yellow, saturate with this color above and below. Costal region of primaries at base a little deeper tinged, and the thorax in front a little ochreous. Under surface of body and legs whitish ; fore tibiae marked with brown ; palpi dark ochreous, white beneath. Fore wings with two stigmata, dark, nearly solid, the outer annulate. Lines blackish, fine, dentate ; the outer line much bent in to below the outer spot, thus differing from *citrina*, and continued on secondaries. No subterminal line. A terminal series of ochrey points on both wings. Fringes pale yellow. Eyes with a white line. Beneath the costa of primaries is shaded with ochreous ; both stigmata repeated ; a common exterior line.

Expanse 19 mil. Texas, Belfrage (No. 383), June and August.

Botis catenulalis, n. s.

Larger than *ventralis*. Entirely brown, not opalescent. Fore wings with the exterior line not greatly indented below vein 2, and followed by a series of dull yellowish spots opposite the scalloping of the line. Discal spots annulate ; all three present ; the inferior (claviform) spherical. The transverse anterior line faintly preceded by a pale shade. Hind wings paler than primaries, especially towards internal margin, crossed by a line corresponding to the outer line of primaries, not much indented, continuous, lunulate, the indentations filled in with pale spots as on fore wings. An even dark brown terminal line ; fringes paler than wing, whereas on primaries these are concolorous. Beneath paler than above with the outer common line distinctly repeated as well as the stigmata on fore wings. Head and palpi brown above, white beneath.

Expanse 26 mil. Hab. California ; Coll. Buff. Soc., from Mr. Meske.

Botis fracturalis Zeller.

Collected by Dr. Shannon in Southern Texas ; also by Belfrage, Nos. 385 and 384 ; varies in color of fore wings as does *argyralis*.
March and April.

Eurycreon communis Grote.

Varies excessively in color. Dark wood brown or fuscous specimens