

(1) For degradation of the cellulose by yeast, 1 g of yeast is added to 10 ml of 2% glucose solution. After 24 hours, the solution is filtered through a Whatman No. 1 filter paper. The filtrate is then used for the experiment.

(2) For degradation of the cellulose by yeast, 1 g of yeast

is added to 10 ml of 2% glucose solution. After 24 hours, the solution is filtered through a Whatman No. 1 filter paper. The filtrate is then used for the experiment.

(3) For degradation of the cellulose by yeast, 1 g of yeast is added to 10 ml of 2% glucose solution. After 24 hours, the solution is filtered through a Whatman No. 1 filter paper. The filtrate is then used for the experiment.

(4) For degradation of the cellulose by yeast, 1 g of yeast is added to 10 ml of 2% glucose solution. After 24 hours, the solution is filtered through a Whatman No. 1 filter paper. The filtrate is then used for the experiment.

(5) For degradation of the cellulose by yeast, 1 g of yeast is added to 10 ml of 2% glucose solution. After 24 hours, the solution is filtered through a Whatman No. 1 filter paper. The filtrate is then used for the experiment.

(6) For degradation of the cellulose by yeast, 1 g of yeast is added to 10 ml of 2% glucose solution. After 24 hours, the solution is filtered through a Whatman No. 1 filter paper. The filtrate is then used for the experiment.

(7) For degradation of the cellulose by yeast, 1 g of yeast is added to 10 ml of 2% glucose solution. After 24 hours, the solution is filtered through a Whatman No. 1 filter paper. The filtrate is then used for the experiment.

30