7.	Arolia free, converging towards the apex. Wing-cell without a
	hook. Pronotum without apical constriction. Genæ very rarely
	high. Loræ sometimes separated also beneath. Eyes inwardly
	mostly emarginate. Tibiæ slender, typically without
	punctures
7a.	Arolia free, converging towards the apex or wanting. Wing-cell
	very rarely without a hook. Pronotum with apical constriction
	(which is sometimes hidden under the posterior margin of the
	posteriorly-produced hind margin of the vertex). Loræ narrow,
-b	sharply separated above and beneath 8, Pilophorini.
70.	Arolia very delicate, or absent (sometimes clearer in some Macrolo-
	phini, but fused with the short claws). [Wing cell without hook.
7.0	Pronotum with apical constriction (e)]
10.	very rarely with hook. Pronotum without apical constriction.
	Vertex wide. Genæ high. Rostrum strong
8.	Head elongate, feebly declivous. Loræ sharply separated above
	and beneath, narrow. Tarsi very slender. Sides of pronotum
	acute, at least posteriorly
8a.	Head vertical9
9.	Head not strongly elongate ventrally. Clypeus not humpily swollen
	in the middle
9a.	Head ventrally strongly elongate. Clypeus humpily swollen in the
	middle. Genæ very high. Neck very short. Tarsi slender, first
	segment long14, Cylapini.
10.	Arolia very delicate, or fused with the short claws. Loræ sharply
	separated above and below,
	narrow
	Arolia absent. Loræ only separated above, arched11, Garganini.
11.	Body generally robust, never constricted in the middle. Loræ
	generally separated above and below, but wide. Tibiæ often
	robust. Membrane with two cells 12, Halticini (Laboparia).
ila.	Body generally narrow, constricted in the middle. Loræ separated

⁽e) Added in MS. by Dr. Reuter.