

which arrested tuberculous disease. Investigations have now been carried out on human patients, and these form the subject of the following observations. It was originally my intention to complete the research, and especially to gain sufficient experience regarding the application of the remedy in practice, and its production on a large scale before publishing anything on the subject; but in spite of all precautions, so many accounts have reached the public, and in such an exaggerated and distorted form, that it seems imperative, in order to prevent false impressions, to give at once a review of the position of the subject at the present stage of the inquiry. It is true that this review can, under these circumstances, be only brief, and must leave open many important questions.

The investigations have been carried on under my direction by Dr. A. Libbertz and Stabsarzt Dr. E. Pfuhl, and are still in progress. Patients were placed at my disposal by Professor Brieger, from his polyclinic; Dr. W. Levy, from his private surgical clinic; Geheimrath Drs. Fantzel and Oberstabsarzt Kohler, from the Charite Hospital; and Geheimrath v. Bergmann, from the surgical clinic of the University. I wish to express my thanks to these gentlemen.

As regards the origin and the preparation of the remedy, I am unable to make any statement, as my research is not yet concluded. I reserve this for a future communication.*

The remedy is a brownish, transparent liquid, which does not require special care to prevent decomposition. For use, this fluid must be more or less diluted, and the dilutions are liable to undergo decomposition if prepared with distilled water. As bacterial growths soon develop in them

they become turbid, and are then unfit for use. To prevent this, the diluted liquid must be sterilized by heat and preserved under a cotton-wool stopper, or, more conveniently, prepared with a one half per cent. solution of phenol.

It would seem, however, that the effect is weakened both by frequent heating and by mixture with phenol solution, and I have therefore always made use of a freshly-prepared solution. Introduced into the stomach the remedy has no effect. In order to obtain a reliable effect it must be injected subcutaneously, and for this purpose we have exclusively used the small syringe suggested by me for bacteriological work. It is furnished with a small India-rubber ball and has no piston. This syringe can easily be kept aseptic by the use of absolute alcohol, and to this we attribute the fact that not a single abscess has been observed in the course of more than a thousand subcutaneous injections.

The place chosen for the injection, after several trials of other places, was the skin of the back between the shoulder-blades and the lumbar region, because here the injection led to the least local reaction—generally none at all, and was almost painless. As regards the effect of the remedy on the human patient, it was clear from the beginning of the research that in one very important particular the human being reacts to the remedy differently from the animal generally used in experiments, namely, the guinea-pig. A new proof for the experimenter of the all-important law that experiment on animals is not conclusive for the human patient proved extraordinarily more sensitive than the guinea-pig. As regards the effect of the remedy, a healthy guinea-pig will bear a subcutaneous injection of 2 cubic centimetres, and even more, of the liquid without being sensibly affected; but in the case of a full-grown healthy man 0.25 cubic centimetre suffices to produce an intense effect. Calculated by the body-weight, one-fifteen-thousandth

* Doctors wishing to make investigations with the remedy at present, can obtain it from Dr. A. Libbertz, Lueneburger Strasse, 28, Berlin, N. W., who has undertaken the preparation of the remedy with my own and Dr. Pfuhl's cooperation, but I must remark that the quantity prepared at present is but small, and that larger quantities will not be obtainable for some weeks.