

quite variable, some individuals corresponding very closely with Mr. Uhler's description of *Jassus divisus*, with which this may prove identical.

*Thamnotettix eburata* n. sp. Form and aspect of *clitellaria* Say, to which it is closely related. Above brown with a large white discal spot, beneath white; venter marked with black and yellow ♂, or whitish ♀. Vertex pale yellow, whitish on the base, slightly suffused with ferrugineous on each side near the eye; a central impressed line extending about half way to the apex. Face whitish, slightly tinged with yellow. Antennæ white, setæ brown. Pronotum and scutellum dark ferrugineous or fuscous, the latter paler with a transverse impressed black line before the apex, behind which the edges are narrowly yellow, and slightly calloused. Hemelytra;—Clavus fuscous, the common disc with a large, obtusely cordate, ivory white spot which is slightly suffused with yellow; behind this spot is a dark area. Costal half of the corium milky white, almost opalescent, tip broadly fuscous; discal half brownish ferrugineous, paler posteriorly, and shading to fuscous along its junction with the white costal area. Beneath and legs white, immaculate; small spines at tip of the tibiae and tarsal joints slightly embrowned; claws black. Venter black, posterior edge and disc of the segments yellow; the latter with a central black line, represented by dots on the three basal segments; convexum yellow; tergum black. Wings milky hyaline, veins faintly brownish. The female differs from the male in being somewhat paler, in having the discal spot of the elytra more yellowish, and the abdomen pale yellow or whitish. In the venation of the elytra and other structural features this species agrees very closely with *clitellaria*. The ♂ has the last ventral segment slightly concave posteriorly; valves obtusely triangular, black; plates white, sparsely covered with long concolorous hairs. The ♀ has the last ventral segment black, with the lateral margins pale; as in *clitellaria* it is deeply incised each side of the middle, leaving a central tongue, which is minutely emarginate at tip; pygofer pale, suffused with brown toward the central groove, and fringed with long pale hairs. Length, ♂ 5 mm.; ♀ 5½–6 mm. A male was swept from grass near a rivulet at Bracebridge; also taken in the vicinity of South Falls.