

The habit of examining the ear in all cases with head symptoms, will sometimes assist materially in clearing up a diagnosis. I once cured a man from the effects of a supposed sun-stroke, by removing inspissated cerumen, who had been treated for two months in a hospital for cerebral disease.

Patients who have once had impacted wax, are apt to suffer again from the same cause, at least I have seen quite a large proportion of cases in persons who have been affected in the same way before. Such may be advised to have their ears syringed with a solution of bicarbonate of soda and water, about once in two months. The removal of the hardened mass is very often a tedious affair. I once spent an hour a day for a week in removing a mass from the ear of a lady patient. In the interim, the best solvents, such as soda, were used. With previous soaking the canal with a warm solution of soda, say a drachm to the half pint, ten minutes will generally suffice to remove the mass. A good india rubber syringe, holding at least four ounces, should be used, and the auditory canal well straightened by holding up the auricle with the left hand, at the same time syringing with the right. The glass syringes are of no use. The stream sent in should be vigorous but steady, and care taken not to eject it with such force as to cause pain or dizziness. There should never be any pain caused in syringing the ear for any purpose. Where pain is produced, syringing will do harm. A thin bowl is held under the ear by the patient. No assistant is needed. No towel need be placed on the patient's neck, for, with careful manipulation, no water will be spilled.

The ear may contain an astonishingly great quantity of hardened ear wax, and an examination should be made very frequently during the course of the syringing to determine when it is all removed. No after-treatment is necessary. If, however, sounds are oppressive, as they often are, after the removal of large quantities of ear wax, a little cotton may be worn in the meatus for a day or two. The membrana tympani always appears reddened immediately after the removal of the cerumen, and then dull. It will be some days before it regains its normal translucency. If the hearing be not improved immediately on removing the wax, the middle ear should be inflated by Politzer's method. The drum is sometimes sunken in temporarily, and one or two passages of air through the Eustachian tube will restore its position as well as the hearing.

Professor Gross recommends the use of a pick for the removal of impacted wax. This does very well as an aid where the wax is very hard. If it be used, the surgeon should have a mirror on his forehead, and never