570/77		
	TANK R	MERVZE
		VILYAA (S)
	HLY JOURNAL DICINE & SUR	DEVOTED TO
nananan yang kananan kananan kananan kananan kananan kananan kanan kananan kananan kananan kananan kananan kana Kananan kananan kanana k	en solenen er er en besterne ander er e	
Vol. xix. H	alifax, Nova Scotia, A	April, 1907. No. 4

All the ferments and soluble constituents of the gastric juice and the amylopsin of the pancreas gland by combination in prescription, at the will of the physician, in an active and agreeable form

'Diazyme' Clycerole4 ouncesEssence of Pepsine, Fairchild4 ouncesM

This mixture, in two teaspoonful doses three times daily, will last the patient ten days.

During this time this mixture will retain the original activity of the several enzymes of the pancreas and gastric juice, namely, amylopsin (pancreatic díastase) pepsin, milk-curdling.

4 c. c. will convert 20 Gm. (300 grs.) starch into sugar in 10 minutes.

4 c. c. will convert 130 Gm. (2,000 grs.) coagulated egg albumen into complete solution in three hours by the U. S. P. test.

4 c. c. will convert 250 c. c. of milk into a soft curd in 5 minutes.

This provides an agreeable, active mixture of the gastric enzymes and pancreatic amylopsin, with the soluble organic and inorganic constituents of the respective gland cells.

It may be taken immediately after meals, or again after an interval of about three hours. It is best taken without dilution, followed by a sip of cold water.

FAIRCHILD BROS. & FOSTER