

Chenopodium Botrys. An interesting sub-excursion was held by the branch, and the members visited an excavation on Sandy Hill at the invitation of Lt.-Col. White, to examine the flora which had appeared subsequent to the carting away of the surface sand. An interesting feature was that, several plants not observed as growing in the immediate locality before, now appeared, and the above named attractive goose-foot was the most conspicuous amongst these.

Cornus sericea was found at Billings' Bridge.

Cornus paniculata. Several nice bushes were observed on the light-house Island above Aylmtr.

Aspidium Goldianum. A new locality for this grand fern was discovered near Kingsmere in the Chelsea Mountains.

Asplenium angustifolium. A pleasing and somewhat novel record has to be made with regard to this beautiful fern. Some 15 years ago a few plants were discovered near Hemlock Lake. When this same locality was visited last autumn the delicate and pale green fronds could be seen in large clumps extending over nearly an acre.

Amongst the new records two of the Orchids are worthy of special mention *Habenaria virescens* was found in abundance at Thurso by Mr. Scott, *Spiranthes Romanzoffiana* a beautiful and highly-scented Ladies Tresses was found by Mr. Scott at Templeton, and on Kettle Island soon afterwards by Mr. Robert B. Whyte.

JAMES FLETCHER,
R. B. WHYTE,
W. SCOTT,
Leaders.

FLORA OTTAWAENSIS.

The following is a list of the additions to the local list discovered since the last report which have not already been recorded :—

Brassica campestris var. *oleifera*, Elgin Street, Sept. 1, W. Scott.

Fragaria Virginiana, Duch, v. *Illinoensis*, Gray, Rockcliffe, May 12, J. M. Macoun.

Dianthus Armeria, L. Hartwell's Locks, July 10, J. Fletcher.

Amarantus blitoides, Wat., Hartwell's Locks, Sept. 3, W. Scott.

Centaurea nigra, L., Thurso, Aug. 7, W. Scott.