

36. Body yellowish-fuscous, purplish or black without a median black line ..... 35  
 Body in great part greenish-yellow with a distinct median black line  
 Habitat—Europe; Massachusetts (Packard); White Mountain; New Hampshire (Mrs. Annie Trumbull Slosson); Long Island (Nathan Banks); Virginia (Wm. D. Richardson); Ithaca, New York ..... *viridis riparia*, Niels
37. Dorsum, except a few yellowish dots, entirely fuscous. The Massachusetts specimens of *Isotoma tricolor*, together with *Isotoma Belfragei*, *purpureascens* and *plumbea*, belong to *viridis*, Bourlet  
 Habitat—Europe; Massachusetts, and Waco, Texas (Packard); Brazos County, Texas (Nathan Banks); Beverly, Massachusetts (A. P. Morse); California (Schott); Ithaca, New York ..... *viridis*, Bourlet  
 Dorsum distinctly marked with yellowish ..... 36
38. Yellowish with a distinctly marked wide transverse black band on each segment. Habitat—Boreal Europe. \**viridis cincta*, Tull. Yellowish, but not with such a band ..... 34
39. Each segment marked at middle with a loop-shaped mark, the sides of the closed end more distant than those of the open end; the open end at the cephalic end of each segment; the closed end sometimes interrupted; the sides of the segments prominently figured with black. Habitat—Boreal  
 Asia ..... \**viridis arctica*, Schott  
 Each segment marked with three closed and united deltoid-shaped marks. Habitat—Agricultural College, Mississippi (H. E. Weed). ..... *viridis delta*, n. var.  
 Species that could not be placed from lack of specimens and of figures of the claws and mucrones:—  
*Isotoma quadrioculata*, Tullb.—“Segmentum tertium abdominis brevius quam quartum, in quo furcula inserta est. Ocelli 4; 2 in utroque latere capitis. Dentes furculae manubris non longiores, recti; mucrones bidenticulati. Long. 1½ mm.” Habitat—Boreal Europe and America. 1872. Tullberg, Sveriges Podurider, p. 48.  
*Isotoma simetaria*, Linn.—“Segmentum tertium abdominis brevius quam quartum, in quo furcula inserta est. Ocelli nulli. Dentes furculae manubrio fere duplo longiores, recti; mucrones bidenticulati. Long. 1 mm. Habitat—Boreal and Central Europe and Boreal America. 1872. Tullberg, Sveriges Podurider, p. 48.

\* Species not seen.