Tongue rather short; maxillary palpi minute; labial palpi simple, slender, slightly overarching the vertex, the third joint about half as long as the second, and pointed; antennae simple, more than half as long as the primaries, with the basal joint short and a little enlarged; scales of the posterior portion of the vertex a little roughened, but not forming a tuft; eyes globose, of moderate size; ocelli, none.

## H. tortricella. N. sp.

Yellowish or straw color; head and hind wings paler, nearly white; there is a minute, pale brownish spot at the end of the disc. Al. ex. 34 inch. Season, May.

### H. albella. N. sp.

White; a minute, indistinct, ochreous spot at the end of the disc, on the forewings, and a very few widely scattered dark brown scales. There is a brownish, ochreous streak on the outer surface of the second joint of the palpi. Al, ev. 14 inch. Season, June, July and August.

### H. canusella. N. sp.

Pale grayish, almost white, with obscure patches of very pale fuscous on the primaries; a small brown spot within the dorsal margin, before the middle; another a little behind it on the fold, and another at the end of the disc. At the beginning of the ciliae is an obscure, narrow, curved, pale fascia, which is very concave towards the base of the wing. Al. ex.

#### GELECHIA.

# G. thoraccalbella. N. sp.

Second joint of the palpi brush-like, the brush spreading and faintly divided; third joint slender, more than half as long as the second.

Palpi whitish, the second joint ochreous brown at the base, and the scales of the brush tipped with ochreous yellow; face, head, and a wide streak from the head to the apex of the thorax, white. Antennae, sides of thorax over the wings, and the primaries brown, the primaries very faintly streaked with whitish towards the apex. Al. cx. 1% inch.

# G. minimaculcila. N. sp.

Second joint of palpi brush-like; third more than half as long as the second.