



THE MARITIME MEDICAL NEWS

A MONTHLY JOURNAL DEVOTED TO
MEDICINE & SURGERY

VOL. XX. HALIFAX, NOVA SCOTIA, FEBRUARY, 1908. No. 2.

Panopepton

ITS TOTAL SOLIDS

The analysis of Panopepton, printed on the label, gives total dry solids 22%, which is really below the actual percentage, as shown by records of assay.

In estimating the practical value of a food for the sick, it is all-important to consider the nature and availability of the solids, in view of the fact that in many cases the ingestion of even a small amount of insoluble matter might be disastrous—cases where the food stuff must be sterile, soluble and wholly diffusible.

The total solids of Panopepton are derived from prime lean beef and whole wheat through the action of the natural digestive principles of the gastric and pancreatic juices under conditions approximating as closely as possible to those of normal bodily digestion, and represent the entire soluble digestible substance of these two foods in a non-coagulable, highly diffusible and wholly absorbable form.

There is no cane sugar in Panopepton—to increase “total solids” and “calories” to the detriment of its general utility.

The solids of Panopepton are actual available food solids, most acceptably and agreeably conveyed in a stable sterile solution.

FAIRCHILD BROS. & FOSTER

New York

ACADEMY OF MEDICINE
TORONTO.