

- Tides**, report by A. S. Christie, 637-700.
 record of observations, 637, 641-672, 694-696.
 stations, geographical position, 637, 694-696.
 electrical self-recording gauge of Lieutenant Greely, 637.
 tide-gauges and methods of observation, 637, 638, 694, 695; cuts.
 bench-marks, lines of levels, homogeneity of series, 637-640.
 lunitidal intervals and heights, 651-682.
 semimensual inequality in intervals and heights, 674-682, 700,
 plate ii, figs. 2, 3.
 parallactic effect upon, 677-679.
 declination effect upon, 679-682.
 diurnal inequality in times and heights, 682-692, 700, plates i, ii,
 fig. 1.
 half-tide level, 651-672, 674.
 effect of variations of atmospheric pressure upon, 692, 693.
 results of harmonic analysis of one year's observations, 696, 697.
 mean sea-level, 697.
 origin of tides in Lady Franklin Bay, 697-700, chart.
 approximate cotidal lines, 700, chart.
- Time**, Washington used except in magnetics, 481.
 observations for, by Sergeant Israel, 481-486, 490-491.
 and longitude, observations for, at Fort Conger, 492-504.
 computation from transit observations, 1881-'82, 720-724, chart.
- Timmia austriaca*, 17.
Torsion, 481.
Tortula icmadophila, 17.
 leucostoma, 17.
 ruralis, 17.
Total intensity, 628.
Trachomedusa, 43.
Transit, Perry (Coast Survey, No. 11), 481, 482.
 observations and computations of time from, 720-724, chart.
- Trichechus rosmaris*, 8, 9.
Trichotropis borealis, 58.
Tringa canutus, 24, 25.
 maritima, 25.
Trisetum subspicatum, 15.
Tritropis aculeata, 50, 52.
 Trout caught in Lake Alexandra, near Fort Conger, 55, plate.
Tunicata, 50.
Turnstone, 25, 26.
- U. S. Coast and Geodetic Survey**, magnetic, tidal, and pendulum reductions by, 477, 637, 701.
 co-operation with Signal Service in furnishing instruments and instructing observers, 479, 480, 701.
- Uria grylle*, 20.
 troile (or *lomvia*), 21.
Urinator lumme, 20.
- Urinator adamsii*, 28.
Ursus maritimus, 1, 2.
- Van Rensselaer Harbor**, annual decrease of west declination at, 631.
 Vasey, G., Dr., botanical specimens identified by, 11.
Vegetation at "Farthest North," 18.
 Vermes, 48, 49, 53, plates ii, iii.
Vertical intensity, 628.
Vesicaria arctica, 12.
Voitia hyperborea, 17.
Vulpes lagopus, 2.
 lagopus fuliginosus, 2, 3.
- Walrus**, 8, 9.
 Washington mean time used except in magnetics, 481.
 Watson, S., botanical specimens identified by, 11.
Weather, observations from St. John's to Fort Conger, 404, 405.
 record from Fort Conger to Camp Clay, 412, 421.
 record at Camp Clay, 433-436.
 charts, simultaneous, March 8, 9, 1883, 317, charts 15, 16.
- Webera acuminata*, 16.
 cruda, 16.
 longicolla, 16.
 sphagnicola, 16.
- Whale**, right or common, 9.
 white, 9.
- Wheat-ear**, 27, 28.
- Wind**, list of charts and tables, 92, 93.
 report, 257-314.
 hourly direction and velocity, August, 1881, to July, 1883, 258-305.
 method and place of observations, 306.
 comparative velocities, 307.
 high winds, 308.
 monthly frequency and velocity, 309.
 record of various velocities, 310.
 monthly and hourly resultants, 312-314.
 annual and diurnal fluctuation of direction, 314, charts 12, 13.
 observations from St. John's to Fort Conger, 404, 405.
 record from Fort Conger to Camp Clay, 412-421.
 record at Camp Clay, 433-436.
 comparative anemometer readings at Dutch Island and Fort Conger, 439, 440.
- Wind vane**, 306.
 Wohlgemuth, E. von, Captain, cited, 20, 168.
 Wolf, 3.
 Woodruff, T. M., Lieutenant, cited on movements of cold waves, 158.
- Xema sabinii*, 22.
- Ziwet**, A., reduction of magnetic observations by, 477, 511.