Ichneumon infucatus, Cress.—Cat. Hym. N. Am., p. 185. One & received by Mr. Fletcher.

ıi-

r.

n

;.

- Ichneumon insolens, Cress.—Taylor, loc. cit.: "One specimen bred from chrysalis of Vanessa antiopa."
- Ichneumon lividulus, *Prov.*—One ? received by Mr. Fletcher, labelled *Ich. grandis*, determined by Mr. Brodie. Seems, from the partially rufous legs, etc., to belong rather to this species.
- Ichneumon longulus, Cress.—Taylor, loc. cit. A specimen so labelled, received by Mr. Fletcher, is, however, only the 3 of cestus, varying a little from typical coloration.
- Ichneumon nuncius, *Cress.* Three &s; also four received by Mr. Fletcher.
- Ichneumon occidentalis, Hargtn.—Type ? in my collection.
- chneumon otiosus, Say.—Taylor, loc. cit.: "My only specimen was unfortunately destroyed during the process of examination."
- chneumon rufiventris, Brullé.—One Q labelled insulens apparently belongs to this species.
- chneumon russatus, *Cress.*—Two \mathfrak{P} s. Type was from V. I. coll., H. Edw.
- chneumon sagus, Cress.—One Q received by Mr. Fletcher.
- chneumon salvus, Cress.—The 3 was described from V. I. coll., H. Edw. chneumon scibilis, Cress.—One A.
- chneumon seminiger, Cress.-Taylor, loc. cit. Not seen.
- chneumon sequax, Cress.—Type Q was from V. I. coll., H. Edw. Taylor (loc. cit.) says: "Very common; one specimen was bred from the chrysalis of a Lycana."
- chneumon Taylori, Hargtn.—Type Q in my collection.
- chneumon vancouverensis, *Prov.*—Type & was from coll. Taylor, who says (loc. cit.), "This fine insect is abundant, and I have bred it in some numbers from the pupa of a Bombyx." Not seen, but answers to description of neutralis, Cr., from Cal.
- thneumon variegatus, Cress.—One of to Mr. Fletcher.
- oplismenus pacificus, *Cress.*—Q & described from V. I. coll, H. Edw. mblyteles hudsonicus, *Cress.*—Two Q s. One of these is a *var*. with the head and thorax above rufous. Mr. Fletcher also has one Q.
- mblyteles nubivagus, Cress.?—One & var.?
- mblyteles perluctuosus, Prov.—One 2.