

results in therapeia that have followed their administration would warrant us in the belief that, as they become better known and more often prescribed, they will become more highly appreciated.

The salts that I have most frequently used are the bromide, iodide and lactate, and I will speak of them in the above order.

Bromide of strontium is a colorless, transparent salt, occurring in hexagonal crystals. It is somewhat deliquescent. The dose is from five grains to one drachm. It is not incompatible with the bromides of the alkalies, and it is soluble in both water and alcohol; it can be administered with all the alcoholic tinctures and most fluid extracts. Its indications for use are those of bromide of potassium, and, while it is a perfect substitute for the potash salt, its prolonged use even in large doses does not seem to produce the untoward results so often noticed in the use of the former salt. The gastric disturbances, the cutaneous eruptions so often noticed in the use of the potash salt, are not seen when the strontium salt is used; again, the depressing and systemic agitation from the prolonged use of the potash, which all have encountered in practice, I have never seen from the strontium bromide.

In cases of epilepsy and other spasmodic neuroses, where the potash salt has been given for a long time, the patient thereby becoming insusceptible to its action, the strontium salt may be substituted with safety and great advantage.

In many diseases of the stomach, the bromide salt will be found of especial benefit. In three obstinate cases of vomiting of pregnancy in which I have prescribed the drug during the past year, two received signal benefit, while in the third case it seemed to have no marked effect upon the vomiting, as the stomach would not retain the remedy; in this case it appeared to have some reflex effect upon the vomiting centre, when given in drachm doses per rectum every six hours, and it was so administered for several days in connection with other treatment.

In one case of hyperæsthesia of the stomach that accompanied and followed ulceration for several weeks after I was satisfied the ulcer had healed, the neuroses promptly yielded to