by the stemach or by the rectum, and it is not necessary to give a larger dose by the latter than by the former method.

M. Claudo Bornard has shown that less chloroform is necessary to produce anasthesia in an animal or a man previously narcotized by morphia, than in one to whom morphia has not been proviously given. I have observed, climically, that there is a similar mutual action between hydrate of chloral and morphia. A fourth of a grain of one of the salts of morphia, followed in an hour by 15 or 20 grs. of hydrate of chloral, will produce a greater anasthetic and hypnotic effect than a largor dose of oither of them given alone. In many cases, where a certain dose of morphia is sufficient to allay pain, but not to produce sleep, it is better to follow the morphia by hydrate of chloral than to repeat the former.

In concluding this account of the physiological and therapeutical action of chloral, I can I express my opinion of it
more clearly than to quote the words of Dr. DaCosta in the
paper previously referred to, viz... "that it is an important
addition to our therapeutic means—chiefly, however, as a hypnotic. Its action as a destroyer of pain is limited; nor do I
think it can compete with other or chloroform to produce the
insensibility requisite for surgical operations. Its chief value
seems to be that of an auxiliary to opium, or to take its place
whon opium is not admissible." As a hypnotic, hydrate of
chloral is better than opium. Chloral sleep is better than opium
steep.—Boston Med. Journal.

TRATHEN OF RESEMBLE FEVER BY PERCHLORIDE OF IRON.—The marked effects of tineture of perchlorade of iron in such discusses as cryv-ipelas and dijhtheroid sore throat had induced J. Russell Reynolds, M.D., F.R.S., London, to try it in acute rheumatism—which agreed with the others in coming under the class of "spreading" inflammatory affections. He had given it in eight cases with such sinces as would justify a further trait. Having given brief histories of the eight cases, he directed attention to certain points. I The relief of the joint affections was definite uniform and speedy. In our cases it was removed in one day, and the longest period of suffering