

EXPLANATION OF PLATE.
Psylla floccosa, new species.

1. Fore wing.
2. Hind wing.
3. Genital segment of female.
- 4, 5, 6. Details of foregoing.
7. Genital segments of male.
- 8, 9, 10. Details of foregoing.
11. Front leg.
12. Second leg.
13. Showing ventral spur at base of third leg.
14. Third leg, showing tibial and tarsal spurs.
15. Head of male.
16. Head of female.
17. Nymphs in flocculent mass on terminal leaves of *Alnus incana*,
June 10, 1909.

NEW PSEUDOSCORPIONIDA.

BY NATHAN BANKS, EAST FALLS CHURCH, VA.

In the CANADIAN ENTOMOLOGIST for 1891, pp. 161-166, and 1893, pp. 64-67, I described a number of these small Arachnids, and in the Journ. N. Y. Ent. Soc., 1895, pp. 1-13, I presented a review of the forms then known from North America. Since then many specimens and species have been collected, and in these pages ten of the new forms are characterized, and a new genus erected for a form near *Garypus*.

Chelifer fuscipes, n. sp.—Cephalothorax and palpi dark brown, fingers more reddish, dorsum of abdomen brown, with a broad pale median stripe, legs brown, paler at tips of some of the joints. Cephalothorax subtriangular, minutely granulate, with a few short, nearly clavate hairs; eyes distinct. Palpi with fine short subclavate hairs, except on fingers, hand and outer side of the tibia, which are simple; trochanters with distinct tubercle behind, femora a little longer than cephalothorax, subpedicellate, largest at middle; tibiæ about as long as femur, pedicellate, evenly convex both sides, but a little broader than femur; hand barely longer than tibiæ, about twice as broad as femur, fingers shorter than hand, stout, curved. Dorsal abdominal scuta finely granulate, outer side very distinctly prolonged behind into acute spines.

Length, 1.8 mm.

From Claremont, Calif. (Baker).

September, 1909