

- Noctua clandestina*—19th June to July ; common ; at sugar and lime blooms.
- brunneicollis*—2nd July to September ; rare ; at sugar.
- alternata*—1st July to September ; very common ; at sugar.
- cupida*—17th July to August ; frequent ; at sugar.
- rubi*?—4th August ; rare ; at sugar.
- Agrotis herilis*—31st July to September ; very common ; at rest, sugar and light.
- tricosa*—3rd August to September ; very common ; at rest, sugar and light.
- subgothica*—9th August to September ; not so frequent as the above two species ; at rest, sugar and light.
- All three species in abundance on flowers of Thistle (*Cirsium arvense*) and unexpanded flowers of *Venibascum thapsus*.
- fennica*—10th August ; one specimen at rest in the Montebello Gardens.
- gladiaria*. *N. sp.* Morr.—Sept. 5th ; not unfrequent at sugar and light.
- tesselata*—29th June ; bred from larva found in abundance at the roots of *Malva rotundifolia* in May ; afterwards to 2nd of July, swarming ; at sugar, rest and at lime blooms. Many remarkable and beautiful varieties.
- Cochrani*—27th July ; bred afterwards ; common ; at sugar and rest to September.
- saucia*—14th August ; not uncommon ; at sugar.
- suffusa*—2nd June to October ; swarming at sugar, flowers and light.
- venerabilis*—6th September ; not uncommon ; at light and sugar ; also bred.
- Aplecta pressa*. *N. sp.*—6th July ; lime blooms and at rest ; several.
- herbida*—27th June to September ; common at rest ; less frequent at sugar.
- nimbosa*—6th August ; rare ; at sugar.
- latex*—30th May to June ; not uncommon ; at rest.
- Ammonoconia badicollis*—31st July ; rare ; at sugar.
- Hadena subjuncta*—2nd July ; rare at sugar.
- vicina*. *N. sp.*—4th June ; rare at sugar.
- confusa*—8th May ; rare at palms.