

***Argynnis aphrodite*, Fabr.**

This butterfly is common throughout the Province. It is frequently mistaken for *A. cybele* which it very much resembles, but may be separated from that species by its smaller size. Flies in July and August.

***Argynnis atlantis*, Edw.**

Common at St. John, Fredericton, Sussex, Belleisle, Moncton and Chipman. This butterfly is more abundant than *A. aphrodite*, frequenting the same localities and flying in company with that species.

***Argynnis myrina*, Cram.**

(*BRETHIS MYRINA*.)

Abundant throughout New Brunswick. This is our most common *Argynnis*, flying from the latter part of June to the last of August.

***Argynnis bellona*, Fabr.**

(*BRETHIS BELLONA*.)

This species is rare in St. John County, and does not appear to be numerous in any part of New Brunswick.

***Melitæa phæton*, Drury.**

(*EUPHYDRYAS PHÆTON*.)

Rare; a few specimens have been captured near St. John. On June 18th of the present year, Mr. Geo. W. Bailey captured a specimen at Springhill, York County.

***Melitæa harrisii*, Send.**

(*CINCLIDIA HARRISII*.)

A specimen of this species was taken near St. John by Philip J. R. McIntosh during the summer of 1897. Rev. C. J. S. Bethune (*The Butterflies of the Eastern Provinces of Canada*) gives New Brunswick as a locality for this rare Canadian butterfly.

***Phyciodes tharos*, Drury.**

Abundant from early in July to September. The form *MARCIA*, Edw. being taken in the earlier part of the season, and the form *MORPHEUS*, Fabr. later. It is found in open meadows and fields during June, July and August.

***Grapta interrogationis*, Fabr.**

(*POLYGONIA INTERROGATIONIS*.)

Forms *UMBROSA*, Lint. *MORPHEUS*, Fabr.

Rare in the vicinity of St. John, and probably not common in any part of New Brunswick.