

## EURYOPIS: Table of Species.

1. Posterior dorsum of abdomen mostly silvery-white.....3.  
Entire dorsum mostly black, only a few small silvery spots, no bands on legs.....2.
2. With four or five silvery spots; legs with black line under femora .....5-*maculata*.  
With six or ten silvery spots; legs without black line .....*argentata*.
3. Tibia and metatarsus brownish, and without bands; abdomen silvery, with narrow median black stripe.....*formosa*.  
Tibia and metatarsus yellowish, banded with black.....4.
4. Cephalothorax and sternum yellowish.....6.  
Cephalothorax and sternum dark brown.....5.
5. Dorsal silvery mark extends around base; marks on legs fine and much broken up.....*scriptipes*.  
Dorsal silvery mark not around base; marks on legs in form of distinct rings .....*funnebris*.
6. Legs heavily marked with fine black lines; venter silvery. *Californica*.  
Legs only slightly marked with black; venter brownish.....*Texana*.

*Euryopsis argentata*, Emer.—This species occurs in the Atlantic Coast States.

*Euryopsis 5-maculata*, Banks.—I have seen this only from the vicinity of Washington, D. C.

*Euryopsis scriptipes*, n. sp.—Cephalothorax a uniform brown, eyes on black spots; abdomen black, with a silvery-white mark around on upper sides, barely or just meeting on base in front, very broad behind, and enclosing a black heart-shaped dorsal mark; sometimes the white is broader and the black is smaller and indented on its sides; venter black in middle, sides silvery, sternum black, coxæ tipped with black, the spinnerets surrounded with black; legs yellowish, densely marked with small transverse black marks, those at apex of joints heavier and forming rings, the tarsi barely, if at all, marked, the femora but little marked above; the tarsus of male palpus black, in the female the other joints marked with black. Eyes and other structure as in *E. funnebris*, and size the same.

Specimens from Colorado (Osler); Pecos and Beulah, New Mexico (Cockerell).