Neumægen's grand collection, and I have figured a good number of the There can thus be but few cases of doubt as to what I have described." All of these collections I have studied, and my references are to identifications made in them. I do not blame Mr. Grote for making errors, and had he not assumed so infallible a standard for himself in his criticism of others, would not have so often pointed them out. I am said also to have followed Mr Grote's synonymy or "adopted" it. Altogether 59 species have synonyms. Of these 23 are originally stated by myself, Mr. Strecker gives one, Mr. Butler is responsible in whole or in part for six, two of them are mere citations of preoccupied names, and of the older species the synonymy is "adopted" by Mr. Grote from Walker in several instances. This is not scientific literature by any means, and I regret being responsible for it, but I cannot allow Mr. Grote's statement "that I have at least laid down the foundation for its proper study" to go unchallenged. Mr. Grote's work in the North American Noctuidæ has been a necessary one, and has been largely drudgery. No one can better appreciate than I the labor involved in identifying material, naming and describing it. That he made synonyms was simply natural and unavoidable, and is in no wise to his discredit. I expect to make them myself, and have done so already. Our noctuids are far from completely known, and in the Agrotes alone will reach nearly 500 species. I know of more than 20 already that are different from any described in the monograph. Mr. Grote's earlier papers were, as a rule, careful and easy to work with, and so up to the period of Dr. Harvey's work. That Mr. Grote really described Dr. Harvey's species has been often told me; but it is interesting to have the statement from headquarters. Mr. Grote's work in the later period failed to equal the earlier papers, so far as value to the student is concerned, from the fact that he assumed in general that his readers knew the Noctuidæ just as well as he did himself. A brief indication, perfectly characteristic in Mr. Grote's view, was absolutely incomprehensible to one not so well grounded. Mr. Grote's work is essentially descriptive, rarely systematic, never monographic. His generalizations are often well put, interesting and valuable; but withal I have not found anywhere any "foundations" for monographic work that did not already exist in literature. Mr. Grote's correction of my reference to insulsa is just. I somehow overlooked the comparison to repentis. A specimen in Dr. Bethune's collection named by Walker, and agreeing with his description, is a species of Hadena, allied to acvastatrix in maculation, but much darker and richer brown in colour, and is Mr. Grote's H. ducta. Walker's determinations are not reliable, and I do not say Mr. Grote is wrong. JOHN B. SMITH.