Chelanops (Chernes).

As shown by Simon Chernes, Menge is Chelanops, Nicolet. A new species from Long Island may be described as follows:—

Chelanops tristis, nov. sp.

Length 2. mm. Colour pale reddish-yellow, soft parts white. Second joint of palpi with two projections behind; third joint pedicellate, nearly cylindrical, twice as long as second; fourth scarcely shorter than the third, pedicellate, inner margin very convex; hand, pedicellate, broad, inner margin very convex, outer but slightly so; fingers as long as hand, curved; whole palpi furnished with only simple hairs. Body with simple hairs and a few somewhat clubbed ones near the anterior part of the cephalothorax. Legs nearly white. Abdomen oblong oval, dorsal scutæ widely separated, the dark spot much nearer the inner than the outer end. On the sea shore of Long Island, New York.

The described species of *Chelanops* may be separated by the following table:—

Palpi with clubbed hairs
Palpi as long as body, large, very pale pallidus. Palpi not as long as body, small, dark sanborni.
Fingers almost one-half shorter than handacuminatus. Fingers as long or scarcely shorter than the hand4
Dark spots in dorsal scutæ much nearer median that lateral end, scutæ widely separated

Garypus, Koch.

In this genus the cephalothorax is quite suddenly narrowed in front of the eyes. It has not been recorded from the U. S. Several specimens of a species of this genus were found at Ithaca, N. Y., in the crevices of a rocky cliff. The form may be described as follows:—

Grypus granulatus, nov. sp.

Length, 1.7 mm.; colour, abdomen yellowish, legs white, palpi and cephalothorax pale reddish. Palpi longer than the body; second joint short, very convex in front; third not very long, short pedicellate, gradually growing thicker; fourth predicellate, shorter than the preceding,