

EXCISION OF THE CERVIX UTERI

IN CASES OF LONG STANDING LACERATION AND OF PROLIFERATING ENDOMETRITIS. (NINETY-ONE CASES.)

BY T. JOHNSON-ALLOWAY, M.D.,
MONTREAL.

Probably the greatest advance made in uterine surgery of late years has been Emmet's trachelorrhaphy for the repair of laceration of the cervix uteri. Before Emmet's genius had worked reform in the treatment of this lesion, afflicted patients were subjected to the enforcement of various blind routine procedures, which we at present do not feel very proud of. But although the key-note of reform in this matter was first struck by Emmet, it is no reason why we should rest content and abstain from effort at still greater improvement in methods for the repair of the lesion in question. Several years ago I recognized the difficulty of obtaining good results by Emmet's method in cases of old standing lesions, where great eversion had existed, and where connective tissue hypertrophy with extensive cystic disease were present. On removal of the tissues at the margins and high up in the angles of the tear, an elevated central plateau of diseased gland and connective tissue structure, situated in the centre of each lip, was left. When these central ridges had been tightly approximated by the application of the sutures, two serious defects became obvious. No provision for drainage after curetting the uterine cavity, and tension so extreme was brought to bear upon the sutures, that in many cases the ones nearest the apex cut out, and allowed a bad ectropium to result. Probably Dr. Emmet does not curette the uterus at the same sitting, and therefore would not require to provide for drainage; he may also introduce his sutures in such a manner that they do not cut out so frequently as in other and less skillful hands. Be this as it may, results were always excellent in recent cases, but were not so in the more chronic.

It occurred to me about this time that the defect may have been in the pathology and that we had not due regard for the diseased ridges left in the centre of each cervical segment, and from this standpoint the operation was really only partially exe-