

the methods introduced by McGraw, that employed by Mikulicz, Moynihan's method, or the method developed by Robson-Murphy's oblong button; or Connel's suture method can be employed in connection with the methods of Mikulicz or Robson, but it seems likely that the button will continue to lose more and more of its old advocates while it is not likely to gain many new ones. This is true, especially, because with it the size of the opening is virtually limited, and there is a distinct objection in the minds of most surgeons against a non-absorbable foreign body.

The one great point in favor of the button is its ability to punch out an opening, and to leave the union between the stomach and the intestine with the slightest possible amount of connective tissue.

In order to be of any practical value this paper must point out some of the dangers to be avoided in surgery of the stomach.

*Unnecessary Traumatism Should be Avoided.*—There is great danger in unnecessary manipulation, because this increases the shock and the tendency to infection.

In all of these cases much can be done to prevent this by making an ample abdominal incision. Much time is frequently occupied in finding the jejunum, resulting in useless handling of viscera. By simply lifting out the transverse colon, and following its mesentery to a point a little to the left of the median line, one can always find the beginning of the jejunum in a few moments.

In gastrectomy and pylorotomy it is possible to reduce the manipulations to a minimum by simply grasping the four main arteries, and also the greater and lesser omenta between these four points, and then excising the intervening portion, which has been grasped by long-jawed forceps, in order to prevent leakage.

There is danger of necrosis of the stomach, if the gastric artery is injured, and of the transverse colon, if the middle colic artery is grasped in clamping the greater omentum.

In making a posterior gastro-enterostomy, there is danger of contraction of the opening in the mesocolon, unless the edges of this are sutured to the stomach.

There is always danger of angulation of the jejunum at its point of attachment to the stomach.

In all stomach operations it is well to have the patient placed in the sitting or semi-sitting posture, within a few hours after