

18. Punctures of abdomen dense.
- Face with black hairs; first recurrent nervure received before the middle of the second submarginal cell; dorsulum with reddish pubescence *Halli*.
- Face with ochreous hairs; first recurrent nervure received beyond the middle of the second submarginal cell; pleura with pale pubescence *solidula*.
- Face with ochreous hairs; first recurrent nervure received before the middle of the second submarginal cell; pleura with pale pubescence *junonia*.
- Face with pale to dark brown hairs; pleura with pale pubescence *compactiscopa*.
- Punctures of abdomen not dense.
- Pubescence of dorsulum ochreous.
- Pleura pale *vicina*.
- Pleura black.
- Abdomen black:
- Dorsulum entirely pale.
- Face with pale pubescence; first recurrent nervure received before the middle of the second submarginal cell *Carlini*.
- Face with black pubescence; first recurrent nervure received beyond the middle of the second submarginal cell.
- Enclosure smooth *neurona*.
- Enclosure partly rugose *pluvialis*.
- Dorsulum with a black band *transnigra*.
- Abdomen blue *Seattlensis*.
19. Clypeus sparsely punctured, especially in the middle.
- Clypeus deeply punctured.
- Abdomen blue, with a greenish cast *Chapmanæ*.
- Abdomen greenish and purplish *Chapmanæ* race.
- Abdomen black.
- Scopa very compact *Pullmani*.
- Scopa loose *longihirtiscopa*.
- Clypeus not deeply punctured *vicinoides*.