

Legs: coxæ and trochanters all mottled brown and white scales; fore femora dark brown with apical white spot, tibiæ dark, 1st and 2nd tarsal joints with tiny basal white spots, remaining joints dark; mid-femora dark with narrow light line ventrally, apex white, tibiæ dark with very small basal white spots, 1st and 2nd tarsal joints with small basal white spots, and a couple of white scales at base of third joint on one leg, the rest dark; hind femora white at base and as a line on cephalic aspect to near the apex, apex white, tibiæ dark with tiny basal white spot, 1st, 2nd, 3rd and 4th tarsal joints dark with broad white basal bands, last joint pure white.

Wing: brown scaled; cells rather long, 1st submarginal distinctly longer and about the same width as 2nd posterior; base of third long vein and mid cross-vein meet and of about equal length, posterior cross-vein a trifle longer and distant more than twice its length. Halteres dark.

Length 5 mm., without proboscis.

Habitat, Cottabato, Mindanao, P. I.

Taken December.

Described from two quite perfect specimens sent by the Surgeon at the Post.

The mesonotum suggests *D. fusca*, Theob., but the third long vein is not carried back, and the leg markings are, of course, quite distinctive.

*Culex? aureopunctis*, n. sp. (female).

Head brown, covered with ochraceous curved and lateral flat scales, many dark brown fork scales, a line of yellow scales around the eyes, a couple of yellow bristles between the eyes and many brown ones projecting forward around the eyes. The head does not seem at all denuded, nevertheless there is a bare median V-shaped space, the base at the vertex. Antennæ dark brown, white banded at the joints, verticels and pubescence dark brown, basal joint brown; palpi dark brown with some pale scales; proboscis apparently has the apical third partly denuded, but the remaining scales show dark brown with a deep ochraceous band near the apical third especially well marked on the ventral side, labellæ dark; eyes brown with red-gold reflections; clypeus brown.

Thorax dark brown; prothoracic lobes with light ochraceous scales and brown bristles; mesonotum covered with dark brown curved scales and a few golden yellow ones which make a broken line from the wing joints to the prothoracic lobes at the junction of mesonotum and pleura, and a faint line on either side of the "bare space" running cephalad