

One grain of the salt, superacidulated with five drops of strong acetic acid, is the proper proportion for admixture with each fluid ounce of water. It is essential that the salt and the acid should be incorporated before the addition of the water, and that the whole should be brought to the boiling point. Superacidulation is necessary on many accounts; it secures perfect solution, increases the decomposing activity of the liquid, and prevents the formation of any carbonate of lead.

As the salts contained in the urine tend to decompose the solution, and lessen its effects on the concretion, the bladder should be evacuated, and washed out with tepid water before the lead fluid is introduced. A double-current caoutchouc catheter is the best for this purpose, as it enables a continuous stream to be employed; and as, on account of its flexibility, it is less liable to irritate the urethra, which should be sedulously avoided. From four to eight fluid ounces of the solution may be thrown into the bladder at a time, and renewed every ten or fifteen minutes, as often as may be deemed proper. By renewing the liquid at short intervals, much greater effect on the calculus is ensured, than when it is allowed to remain longer; for the precipitate formed by decomposition soon envelopes the stone, and puts a stop to further action, until a fresh surface is exposed. Exercise during the retention of the injection increases its effect. Some slight revulsion may be produced by the first introduction of this, or any other fluid, into the bladder; when such is the case, the operation should be remitted for a day or two, and cautiously renewed. The injection may be either warm or cold, as may be most agreeable to the sensations of the patient. Warmth favours the decomposition of the calculus.

If used with proper precautions, I have found that the lead solutions exert a sedative and salutary influence on the lining membrane of the bladder, as they do on external surfaces under inflammation. They also act upon the mucus, which is so abundantly formed in cases of this nature, coagulating it into short curdy flakes, which are easily passed through the urethra.

When the urethra itself is inflamed, or abraded, the injection will be injurious; for the lining membrane of the canal is, I believe, more sensitive than that of the bladder. The introduction therefore of decomponents should be had recourse to, either before lithotomy, or after the urethra has recovered from the effects of the instrument employed, but can never be used, with any prospect of success, where organic disease of the bladder or prostate exists. The injection should not be employed during the internal exhibition of hydrochloric acid, although it may be freely used when nitric acid is administered. When the bladder is not very irritable, a dilute nitric acid injection, alternating with the lead solution, will hasten decomposition.

The two facts established with respect to the lead salts, viz., first, their toleration by the bladder; and, secondly, their chemical action on calculous concretions, induce me to hope that they may become useful agents in the treatment of various other affections in the urinary organs. I have never presumed to imagine they would prove specific solvents for the stone; but, I trust that, where surgical operation is inadmissible, they will be of some avail for relief, if not for cure, by smoothing asperities, and removing the outward phosphatic coating of calculi, so as to bring them within the verge of the crushing forceps; in short, that they may avail for partial, if not entire disintegration. The latter is more likely to happen where layers, composed of the urates or oxalates, are bound together by phosphatic cement. On this species of calculus, they are calculated to act as highly carbonated waters do on those of another description.