

**RHOPALOCERA, Boisd.**

*Family I. PAPILIONIDÆ, Leach.*

*Sub-Family I. PAPILIONIDI, Steph.*

*Genus 1, PAPILIO, Linn.*

1. *P. Asterias, Fab.*, (Black Swallow-tail).—Abundant, June to September.

*Distribution.*—West Indies, S. America, Georgia, Virginia, (Boisd.); Massachusetts, (Harris); New York, (Timmons); Ohio, (Kirtland); Newfoundland, (Gosse); does not occur in the Eastern Townships, at Sorel or Quebec (?)

2. *P. Turnus, Linn.*, (Tiger Swallow-tail).—Common, May to end of July.

*Distribution.*—Virginia, Georgia, Carolina, (Boisd.); Newfoundland to Mexico, (Gosse); Massachusetts, (Harris); New York and Maine, (Emmons); Ohio, (Kirtland); Eastern Townships, Sorel, Rouge District, Quebec and Southern shores of the Gulf.

*Sub-Family II. PIERIDI, Steph.*

*Genus 2, Colias, Fab.*

1. *C. edusa, Fab.*, (Clouded Yellow).—One specimen, September 10th 1856.

*Distribution.*—New York, (Boisd.); (Eu).

2. *C. Philodice, Godt.*, (Clouded Sulphur).—Very abundant, June to October.

*Distribution.*—Throughout North America.

*Genus 3, Pieris, Schrank; Pontia, Fab.*

1. *P. oleracea, Harris*, (Grey-veined White).—Not very numerous, May and June, August and September.

*Distribution.*—Lake Superior, (Agassiz); Wisconsin and Ohio, (Kirtland); Massachusetts, (Harris); Eastern Townships, (Gosse); Upper Canada, Rouge District, Quebec and Southern shores of the Gulf. *P. Casta*, Kirby, a variety of this species occurs in the Hudson's Bay Territories.

2. *P. Protodice, Boisd.*, (see *G. N. and G.* vol. II, p. 347, pl. VI, figs. 3, 4, 5).—Lachine, (Dr. Barnston).

*Distribution.*—New York, (Boisd.); Rochport, Ohio, (Kirtland); Connecticut.

*Family II. HELICONIDÆ, Swainson.*

*Genus 1, Danais, Boisd.; Euploea, Fab.*

1. *D. Archippus, Fab.*, *Plexippus, Cramer*, (Storm Fritillary); Common, June to August.