

The moth from which the above description is taken was bred from a larva taken from an apple tree about fifty miles north of here, May 28th, 1877. It was at that time an inch long, gray, banded transversely with a number of white lines. It moulted June 6th, when all but one of the white lines were replaced by brown, the ground color remaining the same. After feeding a few days longer, it entered the ground and transformed to a chrysalis as above. At this time it was about an inch and a half long. The imago appeared March 27th, 1878.

---

### ON THE EMERGENCE OF LEPIDOPTERA FROM THEIR COCOONS.

BY C. E. WORTHINGTON, IRVING PARK, ILL.

In the years 1856 and 1857 Capt. Thos. Hutton communicated to the London Entomological Society (Trans. v., 85) and to the Journal of the Agri-horticultural Society of India (ix., 167-9) certain observations on the means employed by the imago of *Actias selene* in obtaining exit from its cocoon. In 1857 Messrs Horsfield and Moore in their catalogue of the Lepidoptera in the Indian Museum, quote and endorse Capt. Hutton's observations, and in the course of their remarks indicate indirectly that the same methods are employed by *Antheraea paphia*, an Indian Attacian allied to our *T. polyphemus*.

In these articles the hooks on the wings and the drop of acrid liquid on the head are both noticed and the conclusion arrived at that the means employed are both chemical and mechanical. Capt. Hutton, however, states that the moth discharges this liquid from the mouth and applies it with the brush on the forehead—apparently an error, as the structure of the mouth parts would hardly admit of the secretion of such a liquid, and when secreted it could hardly be conveyed to the forehead.

On reading these notes it occurred to me that I had noticed that examples of *polyphemus* emerging from cocoons from which the top had been removed invariably carried a drop of brown liquid on the frontal tuft, and a little investigation convinced me that the liquid, so far from being secreted by the mouth, was contained in a cell underlying the conspicuous greenish spot on the pupa. This cell is ruptured from the top