	109
4. Tegulæ black; lower part of cheeks covered with hair; axilla spines rather long (Colorado)	11
Tegulæ red 5. Sides of middle of mesothorax very densely punctured; lower part cheeks covered with hair	
part of cheeks bare, bounded behind by a very strong lead a will	er
spines very short	gs
(Arizona)	1.
Face narrower; legs black, with the small joints of tarsi and apices of femora and tibiae dark red; median apical spine of abdomen ver	
long (California)	
7. Inth abdominal segment with small lateral spines, bein of formalling	
(New Mexico)	
and abdominal segment without lateral spines hair of face with	
Jenowish that (Washington Co., Wis . Graenichen) demand Co.	
o. regular chillery pright red : leg red	
Tegulæ black, or dull and dark red	
9. Done apical teeth of andomen long and slander	
apical teeth of abdomen broadened; thorax above without	
man spots	
o. Ventral surface of abdomen densely and very coarsely punctured	ì
(Boulder, Colorado)	
(Falls Church, Va.; Banks)sayi Robertson.	
t. Larger; lower apical teeth of abdomen more widely separated;	
second s. m. receiving recurrent nervures about equally distant from	
base and apex	
r. n. joining second s. m. very near base, very much nearer than	
second r. n. to apex (California)hirsutissima Ckll.	
2. End of abdomen narrow and elongated, with a deep parallel-sided	
and a dove, lower apical feeth very sharp hind anusa and	
sman species (Doulder, Colorado)	
End of audomen proader, the median sulous broads and	
. The strongly enined	
Legs black, or the tarsi red15.	