

- Myoma, transverse colon, descent of, with, 463
 treatment of, 504, 506, 571, 588, 629, 630, 671
 tubal pregnancy with, 342
 tuberculosis of endometrium with, 335
 of Fallopian tubes with, 341, 396
 of uterus associated with, 421
 tuberculous peritonitis associated with, 464
 tubo-ovarian abscess with, 290, 344, 396
 cysts with, 340
 ulceration of vagina with, 440
 umbilical cyst associated with, 462
 hernia associated with, 461, 639
 umbilicus and, 461
 unfolding of cervical glands with, 302
 unrecognized sarcomatous degeneration in, 193
 ureters in cases of, 378-385
 urination and, 376
 uterine cavity and, 305
 polypi associated with, 325
 stones in, 130
 vagina, size of, with, 439
 vaginal, associated with uterine myoma, 234, 440
 cysts with, 439
 hysterectomy for, 629. See also *Vaginal hysterotomy*
 myoma associated with, 440
 myomectomy for, 571. See also *Myomectomy, vaginal*.
 stones in, 440
- Myoma, varieties of, 1
 vein stones in vagina with, 440
 vesical adhesions with, 366
 calculus with, 375
 vomiting associated with, 447
 weakness with, 457
 weight of, 119, 512
- Myomatous tendency, 4
- Myomectomy, abdominal, 506
 abscess in broad ligament after, 553
 age of patients in cases of, 506, 518
 and hysterectomy on account of development of other myomata, 561
 bed-sores after, 555
 bronchitis after, 554
 bronchopneumonia after, 555
 carcinoma of uterus not discovered during, 416
 cystitis after, 549
 death after, 532, 533, 537, 538, 540, 541, 543, 544, 568
 during pregnancy, 528, 529, 538
 evacuation of pelvic abscess after, 559
 for adenomyoma of uterus, 511
 Myomectomy, abdominal, for cervical myoma, 510
 for cystic myoma, 512
 for interstitial myoma, 508, 521
 for myoma of broad ligament, 510
 for pedunculated myoma, 521
 for submucous myoma, 509, 523
 for subperitoneal pedunculated myoma, 508
 for subvesical myoma, 510
 for suppurating subperitoneal myoma, fecal fistula after, 142
 foul discharge from uterus after, 550
 hemoglobin, low, as contraindication to, 517
 hemorrhage after, 548
 hysterectomy after, 559, 560
 hysterical manifestations after, 547
 results after, 568, 569, 570, 672
 in syphilitic patient, 518
 intestinal adhesions after, 566
 obstruction after, 533, 552, 557
 leukocytosis after, 553
 lobar pneumonia after, 555
 location of myoma and, 507
 myocarditis with, 544
 nausea after, 548
 nephritis after, 550
 number of myomata removed by, 517
 operations on tubes and ovaries during, 526
 performed in conjunction with, 526
 subsequent to, 556, 570
 pain after, 549
 peritonitis after, 533, 552
 phlebitis after, 555
 pleurisy after, 554
 postoperative complications and, 544
 pregnancy following, 566
 preservation of portion of fundus in, 525
 pulse after, 545, 546
 pus from urethra after, 550
 release of adhesions after, 556
 removal of ovarian cysts during, 527
 of ovaries after, 558
 of tubes or ovaries during, 526
 of vermiform appendix and, 463, 557, 558
 repair of perineum during, 527
 repeated, 558
 size of myomata removed, 511
 sloughing of uterine wall after, 551
 suspension of uterus during, 528
 syphilis as contraindication to, 518
 tearing of uterine artery during, 525
 technic of, 521, 523
 temperature after, 545, 546
 tonsillitis after, 554
 vomiting after, 548