

*Salebria levigatella*, n. sp.—Expands 28 mm. Head and thorax blackish. Forewings smooth, of an even fuscous, the costa more rounded than usual, basal space to basal line of a dull red colour, lines indistinct, the basal faintly discernible, dentate, the outer very faint or entirely obsolete. Hindwings fuscous, darker outwardly. Amherst, Mass., from Dr. C. H. Fernald; also from Wisconsin.

*Salebria purpurella*, n. sp.—Expands 27 mm. Face and palpi gray, mixed with black; the summit of head stained with reddish. Thorax and abdomen fuscous, with reddish stain. Forewings gray, heavily washed with reddish-purple, the gray being more clear before basal line and along costa before outer line. Basal line near base, apparent only on posterior half, lined on each side with black; outer line very faint and indistinct close to margin. Hindwings fuscous, tinged slightly with reddish near anterior angle. New Mexico.

*Zophodia bella*, n. sp.—Expands 28 mm. Head and palpi fuscous-gray; thorax fuscous; abdomen fuscous; the segments lighter posteriorly. Forewings gray, clear along costa, washed with fuscous behind subcostal vein, this running in longitudinal lines on the veins and interspaces, being especially distinct on the veins. Basal line suggested by diffuse blackish spots; outer line quite indistinct, oblique from costa, then strongly dentated; a subterminal line of diffuse black spots; two discal dots black, diffuse, the anterior more distinct. Hindwings light fuscous with marginal black line. Massachusetts.

*Ocala*, n. gen.—Labial palpi long, porrect; maxillary palpi distinct, pencil tufted; tongue long, antennæ bent above base with tuft of scales in bend; forewings 11 veins, 4 and 5 stemmed, 10 separate; hindwings 7 veins, 3 and 4 stemmed, 5 wanting. Abdomen in ♂ tufted. Very near to *Dolichorrhinia*, Rag., and differing principally in the presence of the pencil tufted maxillary palpi and tufted abdomen.

*Ocala dryadella*, n. sp.—Expands 18 mm. Palpi and head fuscous gray, thorax light fuscous or dirty white, abdomen same colour with heavy anal tufts beneath and on sides in male; forewings fuscous gray, basal line well out near middle of wing, faint, edged outwardly with diffuse broken black, outer line close to margin, indistinct, rounded, serrated; two black discal dots, very small; a comparatively large, oval black spot on basal space near basal line just above inner margin. Hindwings translucent, light fuscous.