

**Cauliflower**—Early Snowball, Whitehead, Gilt Edge. Cultivate the same as cabbage. Tie up the leaves around the top as soon as the flower is commencing to form. This facilitates blanching and ensures white heads.

**Carrots**—Half Long Chantenay, Danvers and Scarlet Nantes. Sow the seed in clean ground half an inch deep in rows one foot apart from about the 5th of May up until the 10th of June. Thin the first time, as soon as large enough to handle, to two inches apart, second time taking out every other one. In this way you utilize the thinnings for home use.

**Celery**—Paris Golden, White Plume, Giant White and New Rose. Sow in hot beds March 25th to April 1st, covering seeds an eighth of an inch deep. Keep moist. Thin out or transplant into boxes, two by two and a half inches apart. Transplant into open ground 24th of May. Care should be taken to set the plants just so deep as not to cover the crown. When planting, the earth should be well firmed around the roots. Set the rows three feet apart and bleach by placing boards ten or twelve inches wide on each side. By using the boards this crop can be grown closer than where earth is used for banking.

**Corn**—Golden Bantam, Extra Early White Cob Cory, Early Malcolin and Improved Squaw. Sow 15th of May to June 1st. For early planting the seed should not be covered more than one inch deep, while the latest sowings may be sown at a depth of from two to three inches according to the ground and weather conditions. The rows should be three feet apart for the first and last named and four feet for the "Cory," placing six to eight seeds every two feet, and thinning to four plants in the hill. Hoe the soil slightly up to the plant.

**Citrons**—Red Seeded, Colorado Preserving. Sow May 15th one and one half inches deep in rows of three feet apart. This seed takes a long time in germinating. It is more hardy and requires to be sown earlier than other members of the melon family. Sow seeds six inches apart and when rough leaf appears thin to two feet in the row.

**Cucumbers**—Davis Perfect, Improved White Spine, Extra Early Long Green. Sow in the open ground May 24th to June 10th in rows three and a half feet apart, one inch to one and one half inches deep. Drop seeds every two inches and thin to half a foot.

**Lettuce**—(Leaf), Grand Rapids, Black Seeded Simpson; (Head) Toronto Gem, Big Boston, Nonpareil, Three X Solid Head. Sow in hot bed, 10th April, or in open ground as soon as land can be worked. Sowings can be made or plants set between cabbage and cauliflower rows, the crop being cut and out of the way before the land between the rows of the latter crops is required. Sow seed half an inch deep in rows one foot apart. Thin early and leave plants eight inches to one foot apart in the row.

**Onions**—Extra Early Flat Red, Yellow Globe Danvers, Early Flat Danvers. For transplanting Prizetaker and Southport Globe—red, white, and yellow. As soon as the land is in fit condition to work sow one inch deep in rows twelve inches apart, after giving the ground a thorough raking, and if the land is weedy wait a few days before sowing after the first stirring. Then rake it again and sow. This will kill the weeds which have started and give the seed a better chance. As soon as the plants commence to break the ground, run the rake very lightly across the rows. Hoe close up to the row as soon as you can see the plants. Pull all weeds as soon as possible. Where the land is dirty and there is danger of weeds smothering the crop, you can use the new