the inner margin, covered with long hairs as usual, the rest reticulated; of the same color as the fore wings.

Head gray, darker in the male; colar slightly more buff tinted than the ground color of the wings, making a light drab, without marks; thorax gray, a mixture of the ground color of the wings and black, a black line round the posterior margin preceded by a space devoid of black; abdomen of a lighter gray than the thorax. Beneath the thorax and abdomen concolorous with the same parts above, the legs ringed with light.

This species is nearest to Cossus Angrezi Bailey, but differs from that in several particulars. It lacks the dark shading on the fore wings, in the shape and distinctness of the transverse median line on fore wings, and in the shape and position of the subterminal line. The fore wings are a little less rounded at the apices, otherwise they are shaped alike. The hind wings in this species are reticulated over the terminal two-thirds, while in Angrezi the reticulations do not cover quite half. In Angrezi the thorax is black, with the edge of the tegulæ shaded with yellowish gray; in this the thorax and tegulae are both gray. The antennæ are pectinated in both sexes, most heavily so in the male. The under side of the wings repeat the markings of the upper side, but a little blurred. The space between the two transverse lines is a little shaded with brownish. The female has the ovipositor exserted about .30 of an inch.

Described from two examples, one of each sex, taken by David Bruce, in Colorado, in 1887, and in the name dedicated to him.

Tolype distincta, nov. spec.

Expanse, male 1.121/2, the female 1.50 inches.

Male.—In color and markings between *Velleda* and *Laricis*, rather a dark leaden gray; the fore wings crossed by two geminate lines and a subterminal line, similar to those of *Velleda*, but the subterminal line is also inclined to be geminate. These lines are lighter than the ground colour, but not white as in *Velleda*. The geminate lines bordered each side by a shade line that is slightly brownish; fringes concolorous, a whitish gray line at the base. Hind wings nearly the color of fore wings, a pale shade through the middle and at the outer border; fringe like that on fore wings.