

*Pachybrachys Coloradensis*, nov. sp.—Small, above pale grayish-white, form elongate, slender, elytra parallel, striate, punctate. Length,  $1\frac{3}{4}$ – $2\frac{1}{2}$  mm.

Head large, front nearly flat, rather thickly, brownly punctate, with narrow brownish frontal line, and more or less of a cloud on the vertex; eyes distant, antennæ with first five joints and base of sixth light, after that dark, a little more than half the length of the body; thorax thickly and rather evenly brown punctate, except the edges, depressed behind and often with a small brown spot in the depression before the scutel, edge very lightly and indefinitely rounded; elytra cylinder-shaped, covered with striæ of dark brown punctures, which are diffused in the vicinity of the scutellum and slightly irregular behind and below the humerus; lobe moderate, ♀ with a very well marked row of marginal punctures from upper edge of the curve to end of sinuation, ♂ usually not quite as numerous nor well marked, below black, with epimera, sides of abdomen, venter and pygidium whitish-yellow; legs yellow, with darker spots or clouds on thighs; in some specimens the black of the body under the elytra seems to show through the elytra; the sutural row of punctures is on the edge, and gives the appearance of a very fine black line.

Seven ♂'s, five ♀'s, Colorado Springs, Colorado; Coolidge, New Mexico; Clear Lake, Utah. All coll. by Wickham. Type coll., Bowditch.

Of about the size and shape of *P. densus*, Bow., but lighter in colour and at once separated by the punctate striate elytra.

*Pachybrachys densus*, nov. sp.—Small, above pale testaceous, or dirty-yellow, very thickly and finely, diffusely, brown punctate. Length,  $1\frac{3}{4}$ – $2\frac{1}{2}$  mm.

Head slightly convex in front, the usual dark frontal markings hardly visible, thickly and finely punctulate; eyes distant; antennæ dirty-brown, reaching to first segment of abdomen in ♀; thorax wider than long, thickly and evenly punctate without darker clouds, depressed behind, sides subangulate; elytra of same colour as thorax and entirely, diffusely, thickly punctate, though in one specimen the course of the third, fifth and humeral intervals can be traced; lobe small, wholly occupied by punctures, marginal stria very lightly curved and barely sinuate behind; body beneath dark, with the epimera, abdominal sides and pygidium picked out in lighter colour; legs light, with dark rings on thighs.

♂, ♀, Arizona. Type coll., Bowditch.