

Antennæ inserted somewhat below the middle of the face.

Parapsidal furrows not complete. Mandibles 3- and 4-dentate.

1. Fore-wings with a large, smoky area under the whole of the marginal vein. Propodeum with a large neck, tricarinate. Abdomen produced beneath, with a very short petiole. Pedicel shorter than funicle 1, the latter twice longer than wide. Coxæ and femora concolorous, the middle tibia infuscated, the caudal so at base. Scutellum without a cross-suture before apex (or an indicated one). Clypeus sinuate.....*borrowi*, n. sp.

2. Fore wings hyaline.

Ovipositor not extruded. Coxæ above metallic, the legs lemon yellow, the antennæ yellow-brown. Clypeus sinuate rather deeply at apex. Antennæ inserted but a little above the ventral ends of the eyes. Spiracles at the base of a broad hollow, bounded by the lateral carina and a carina laterad of the spiracle, no sulcus. Propodeum tricarinate. Pedicel and funicle 1 subequal.....*onerati* (Fitch).

Coxæ concolorous, the femora brown or washed with metallic or metallic. Scape yellow.

Clypeus distinctly concave or sinuate at distal margin.

Propodeum tricarinate and with a very short neck (irregular rugæ between the carinæ). Funicle 1 somewhat over twice longer than wide, twice the length of the pedicel, 6 a half longer than wide. Tibiæ white. Marginal vein nearly twice the length of the stigmal. Femora usually washed. Funicle and club

black.....*rhodobæini* Ashmead (= *languriæ* Ashmead).

The same but the femora usually deep metallic, the tibiæ golden yellow, the funicle joints somewhat shorter, the flagellum brown...*phycidis* Ashmead (= *piercei* Crawford).

The same as *phycidis* but the tibiæ brown, white at tip, no median carina on propodeum, the foveæ of spiracular sulcus minute.....*cerealellæ* (Ashmead)